

Anti-TNFRSF7 / CD27 Reference Antibody (Organon patent anti-CD27)

Recombinant Antibody Catalog # APR11057

Specification

Anti-TNFRSF7 / CD27 Reference Antibody (Organon patent anti-CD27) - Product Information

Application FC, Kinetics, Animal Model

Primary Accession
Reactivity
Clonality
Isotype
Calculated MW

P26842
Human
Monoclonal
IgG2SA
145 KDa

Anti-TNFRSF7 / CD27 Reference Antibody (Organon patent anti-CD27) - Additional Information

Target/Specificity
TNFRSF7 / CD27

Endotoxin

< 0.001EU/ μg, determined by LAL method.

Conjugation Unconjugated

Expression system

CHO Cell

Format

Purified monoclonal antibody supplied in PBS, pH6.0, without preservative. This antibody is purified through a protein A column.

Anti-TNFRSF7 / CD27 Reference Antibody (Organon patent anti-CD27) - Protein Information

Name CD27 (HGNC:11922)

Function

Costimulatory immune-checkpoint receptor expressed at the surface of T-cells, NK-cells and B-cells which binds to and is activated by its ligand CD70/CD27L expressed by B-cells (PubMed:28011863). The CD70-CD27 signaling pathway mediates antigen- specific T-cell activation and expansion which in turn provides immune surveillance of B-cells (PubMed:<a href="http://www.uniprot.org/citations/28011863" target="http://www.uniprot.org/citations/28011863" target="http://www.uniprot.org/citations/28

href="http://www.uniprot.org/citations/28011863" target="_blank">28011863). Mechanistically, CD70 ligation activates the TRAF2-PTPN6 axis that subsequently inhibits LCK phosphorylation to promote phenotypic and transcriptional adaptations of T-cell memory (PubMed:38354704).



In addition, activation by CD70 on early progenitor cells provides a negative feedback signal to leukocyte differentiation during immune activation and thus modulates hematopoiesis (By similarity). Negatively regulates the function of Th2 lymphocytes in the adipose tissue (By similarity).

Cellular Location

Cell membrane; Single-pass type I membrane protein

Tissue Location


Found in most T-lymphocytes.

Anti-TNFRSF7 / CD27 Reference Antibody (Organon patent anti-CD27) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture


Anti-TNFRSF7 / CD27 Reference Antibody (Organon patent anti-CD27) - Images



Anti-TNFRSF7 / CD27 Reference Antibody (Organon patent anti-CD27) on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%



The purity of Anti-TNFRSF7 / CD27 Reference Antibody (Organon patent anti-CD27)is more than





95% ,determined by SEC-HPLC.