

**MYD88 Antibody**  
**Catalog # ASC10065****Specification**

---

**MYD88 Antibody - Product Information**

Application	WB, E
Primary Accession	<a href="#">U70451</a>
Other Accession	<a href="#">U70451</a> , <a href="#">1763090</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	Aprox: 35 kDa KDa
Application Notes	MYD88 antibody can be used for detection of MyD88 by Western blot 0.5 µg/mL. For immunohistochemistry use a concentration of 4 µg/mL. For immunofluorescence and immunocytochemistry use a concentration of 5 µg/mL.

**MYD88 Antibody - Additional Information**

Gene ID	4615
<b>Other Names</b>	
MYD88 Antibody: MYD88D, myeloid differentiation primary response gene (88)	

**Target/Specificity**  
MYD88;**Reconstitution & Storage**

MYD88 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

**Precautions**

MYD88 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.


**MYD88 Antibody - Protein Information****MYD88 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **MYD88 Antibody - Images**



Western blot analysis of MyD88 in Jurkat whole cell lysate with MyD88 antibody at 0.5 µg/mL.

### **MYD88 Antibody - Background**

**MYD88 Antibody:** The pro-inflammatory cytokine IL-1 induced cellular response requires IL-1 receptor complex including IL-1RI and IL-1RAcP. Recently, MyD88 was identified as an adapter molecule in the IL-1 signaling pathway. MyD88 associates with and recruits IRAK to the IL-1 receptor complex in response to IL-1 treatment and dominant negative form of MyD88 attenuates IL-1R-mediated NF-κB activation. MyD88 is also employed as a regulator molecule by IL-18 receptor and human Toll receptor, which are members in the Toll/IL-1R family of receptors. Targeted disruption of the MyD88 gene results in loss of cellular responses to IL-1 and IL-18, and MyD88-deficient mice lack responses to bacterial product LPS that employs Toll-like receptors 2 and 4 (TLR2 and TLR4) as the signaling receptors. MyD88 is a general adapter protein for the Toll/IL-1R family of receptors and plays an important role in the inflammatory response induced by cytokines IL-1 and IL-18 and endotoxin. MyD88 gene is expressed in many tissues.

### **MYD88 Antibody - References**

- Muzio M, Ni J, Feng P, Dixit VM. IRAK (Pelle) family member IRAK-2 and MyD88 as proximal mediators of IL-1 signaling. *Science* 1997;278:1612-5
- Adachi O, Kawai T, Takeda K, Matsumoto M, Tsutsui H, Sakagami M, Nakanishi K, Akira S. Targeted disruption of the MyD88 gene results in loss of IL-1- and IL-18-mediated function. *Immunity* 1998;9:143-50
- Medzhitov R, Preston-Hurlburt P, Kopp E, Stadlen A, Chen C, Ghosh S, Janeway CA Jr. MyD88 is an adaptor protein in the hToll/IL-1 receptor family signaling pathways. *Mol Cell* 1998;2:253-8
- Kawai T, Adachi O, Ogawa T, Takeda K, Akira S. Unresponsiveness of MyD88-deficient mice to endotoxin. *Immunity* 1999;11:115-22