

# DcR3 Antibody

Catalog # ASC10070

## Specification

# **DcR3 Antibody - Product Information**

Application Primary Accession Other Accession Reactivity Host Clonality Isotype Calculated MW Application Notes

WB, IHC-P, E <u>O95407</u> <u>AF104419</u>, <u>4106877</u> Human, Mouse, Rat Rabbit Polyclonal IgG 33 kDa KDa DcR3 antibody can be used for detection of DcR3 expression by Western blot 0.5 μg/mL. An approximately 33 kDa band can be detected. Antibody can also be used for immunohistochemistry starting at 1 μg/mL.

## DcR3 Antibody - Additional Information

Gene ID

**Other Names** 

8771

DcR3 Antibody: M68, TR6, DCR3, M68E, DJ583P15.1.1, UNQ186/PRO212, Tumor necrosis factor receptor superfamily member 6B, Decoy receptor 3, DcR3, tumor necrosis factor receptor superfamily, member 6b, decoy

Target/Specificity TNFRSF6B;

### **Reconstitution & Storage**

DcR3 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

### Precautions

DcR3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

# **DcR3 Antibody - Protein Information**

Name TNFRSF6B

Synonyms DCR3, TR6

Function

Decoy receptor that can neutralize the cytotoxic ligands TNFS14/LIGHT, TNFSF15 and TNFSF6/FASL. Protects against apoptosis.



**Cellular Location** Secreted.

## **Tissue Location**

Detected in fetal lung, brain and liver. Detected in adult stomach, spinal cord, lymph node, trachea, spleen, colon and lung. Highly expressed in several primary tumors from colon, stomach, rectum, esophagus and in SW480 colon carcinoma cells

### **DcR3 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

#### **DcR3 Antibody - Images**



Western blot analysis of DcR3 in human heart (H), brain (B), and kidney (K) tissue lysates with DcR3 antibody at 1:500 dilution.



Immunohistochemistry of DcR3 in human heart tissue with DcR3 antibody at 1  $\mu$ g/mL. **DcR3 Antibody - Background** 



DcR3 Antibody: Apoptosis is induced by certain cytokines including TNF and Fas ligand in the TNF family through their death domain containing receptors. Several novel members in the TNFR family including DR3, DR4, DR5, and DR6 were recently discovered and function as cell death receptors. Two decoy receptors, DcR1 and DcR2, were recently identified to compete with DR4 and DR5 for their ligand TRAIL binding. A novel decoy receptor was more recently discovered and designated DcR3 and TR6, respectively,. Unlike DcR1 and DcR2, DcR3 is a soluble rather than a membrane associated molecule. DcR3 binds to FasL and LIGHT and inhibits FasL and LIGHT induced apoptosis. DcR3 transcript is expressed in a number of lung and colon carcinomas and in some normal tissues.

## DcR3 Antibody - References

Pitti RM, Marsters SA, Lawrence DA, Roy M, Kischkel FC, Dowd P, Huang A, Donahue CJ, Sherwood SW, Baldwin DT, Godowski PJ, Wood WI, Gurney AL, Hillan KJ, Cohen RL, Goddard AD, Botstein D, Ashkenazi A. Genomic amplification of a decoy receptor for Fas ligand in lung and colon cancer. Nature 1998;396:699-703

Yu KY, Kwon B, Ni J, Zhai Y, Ebner R, Kwon BS. A newly identified member of tumor necrosis factor receptor superfamily (TR6) suppresses LIGHT-mediated apoptosis. J Biol Chem 1999;274:13733-6 (RD1299)