

**DRAK2 Antibody**  
**Catalog # ASC10073****Specification**

---

**DRAK2 Antibody - Product Information**

Application	WB, ICC, E
Primary Accession	<a href="#">O94768</a>
Other Accession	<a href="#">AB011421</a> , <a href="#">3834355</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	DRAK2 antibody can be used for detection of DRAK2 by Western blot 0.5 µg/mL. An approximately 45 kDa band can be detected. Antibody can also be used for immunocytochemistry starting at 10 µg/mL.

**DRAK2 Antibody - Additional Information**Gene ID **9262****Other Names**

DRAK2 Antibody: DRAK2, DRAK2, DAP kinase-related apoptosis-inducing protein kinase 2, serine/threonine kinase 17b

**Target/Specificity**

STK17B; It has no cross responses to DAP or ZIP kinases. The approximately 70 kDa band is probably non-related to DRAK2 although it is peptide blockable.

**Reconstitution & Storage**

DRAK2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

**Precautions**

DRAK2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**DRAK2 Antibody - Protein Information****Name** STK17B**Synonyms** DRAK2**Function**

Phosphorylates myosin light chains (By similarity). Acts as a positive regulator of apoptosis.

**Cellular Location**

Nucleus. Cell membrane. Endoplasmic reticulum-Golgi intermediate compartment.  
Note=Colocalizes with STK17B at the plasma membrane.

#### Tissue Location


Highly expressed in placenta, lung, pancreas. Lower levels in heart, brain, liver, skeletal muscle and kidney

#### DRAK2 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### DRAK2 Antibody - Images



Western blot analysis of DRAK2 in Raji cell lysate with DRAK2 antibody at (A) 1 and B (2)  $\mu\text{g/mL}$ .



Immunocytochemistry of DRAK2 in Jurkat cells with DRAK2 antibody at 10  $\mu\text{g/mL}$ .

#### DRAK2 Antibody - Background

DRAK2 Antibody: Apoptosis is mediated by death domain containing adapter molecules and a

caspase family of proteases. Certain serine/threonine protein kinases, such as ASK-1 and RIP, are mediators of apoptosis. Two novel serine/threonine kinases that induce apoptosis were recently identified and designated DRAK1 and DRAK2 (for DAP kinase-related apoptosis-inducing protein kinases). DRAKs contain an N-terminal kinase domain and a C-terminal regulation domain. Overexpression of DRAK2 induces apoptosis. DRAKs have high sequence homology to DAP and ZIP kinases, and they represent a novel family of serine/threonine kinases, which mediates apoptosis through their catalytic activities. DRAK2 is located in nucleus and the messenger RNA was ubiquitously expressed in human tissues.

#### **DRAK2 Antibody - References**

Sanjo H, Kawai T, Akira S. DRAKs, novel serine/threonine kinases related to death-associated protein kinase that trigger apoptosis. J Biol Chem 1998;273:29066-71 (RD1299)