

BCMA Antibody
Catalog # ASC10131**Specification**

BCMA Antibody - Product Information

Application	WB, IHC-P, IF, E
Primary Accession	Q02223
Other Accession	NP_001183 , 23238192
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	Predicted: 20 kDa

Application Notes	Observed: 27 kDa KDa BCMA antibody can be used for detection of BCMA by Western blot at 5 µg/mL. Antibody can also be used for immunohistochemistry starting at 10 µg/mL. For immunofluorescence start at 20 µg/mL.
-------------------	---

BCMA Antibody - Additional Information

Gene ID **608**

Other Names

BCMA Antibody: BCM, BCMA, CD269, TNFRSF13A, BCM, Tumor necrosis factor receptor superfamily member 17, B-cell maturation protein, tumor necrosis factor receptor superfamily, member 17

Target/Specificity

TNFRSF17;

Reconstitution & Storage

Antibody can be stored at 4°C up to one year. Antibodies should not be exposed to prolonged high temperatures.

Precautions

BCMA Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

BCMA Antibody - Protein Information

Name TNFRSF17

Synonyms BCM, BCMA

Function

Receptor for TNFSF13B/BLyS/BAFF and TNFSF13/APRIL. Promotes B-cell survival and plays a role in the regulation of humoral immunity. Activates NF-kappa-B and JNK.

Cellular Location

Cell membrane; Single-pass type III membrane protein. Endomembrane system; Single-pass type III membrane protein Note=Perinuclear Golgi-like structures


Tissue Location

Expressed in mature B-cells, but not in T-cells or monocytes


BCMA Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BCMA Antibody - Images

Western blot analysis of BCMA in (A) human spleen tissue lysate, (B) K562, and (C) U937 cell lysates with BCMA antibody at 5 µg/mL.



Immunohistochemistry of BCMA in human spleen tissue with BCMA antibody at 10 µg/mL.



Immunofluorescence of BCMA in Human Spleen tissue with BCMA antibody at 10 µg/mL.

BCMA Antibody - Background

BCMA Antibody: Members in the TNF superfamily regulate immune responses and induce apoptosis. Two novel members in the TNF family were recently identified and designated BAFF/BLyS/TALL-1/THANK/zTNF4 and April/TALL-2, respectively. BAFF was characterized as a B cell activator since it induced B cell proliferation and immunoglobulin secretion. April regulates immunological and non-immunological cell growth. Three receptors, BCMA (for B cell maturation protein), TACI, and BAFF-R, for BAFF were recently identified. BCMA, like TACI, binds BAFF and April. BAFF and its receptors induce B cell development and survival, activate NF-κB and c-jun N-terminal kinase, and are involved in B cell associated autoimmune diseases.

BCMA Antibody - References

Moore PA, Belvedere O, Orr A, et al. BLyS: member of the tumor necrosis factor family and B lymphocyte stimulator. *Science* 1999; 285:260-3
Schneider P, MacKay F, Steiner V, et al. BAFF, a novel ligand of the tumor necrosis factor family, stimulates B cell growth. *J. Exp. Med.* 1999; 189:1747-56
Shu HB, Hu WH, Johnson H. TALL-1 is a novel member of the TNF family that is down-regulated by mitogens. *J. Leukoc. Biol.* 1999; 65:680-3
Mukhopadhyay A, Ni J, Zhai Y, et al. Identification and characterization of a novel cytokine, THANK, a TNF homologue that activates apoptosis, nuclear factor-κB, and c-Jun NH2-terminal kinase. *J. Biol. Chem.* 1999; 274:15978-81