

Anthrax Edema Factor Antibody
Catalog # ASC10279**Specification**

Anthrax Edema Factor Antibody - Product Information

Application	E
Primary Accession	P40136
Other Accession	P40136 , 729244
Reactivity	Bacteria
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	Anthrax Edema Factor antibody can be used for the detection of Anthrax PA protein in ELISA. It will detect 10 ng of free peptide at 1 µg/mL.

Anthrax Edema Factor Antibody - Additional InformationGene ID **3361726****Other Names**

Anthrax Edema Factor Antibody: Calmodulin-sensitive adenylate cyclase, ATP pyrophosphate-lyase, EF, Calmodulin-sensitive adenylate cyclase

Target/Specificity

pxo1_122;

Reconstitution & Storage

Anthrax Edema Factor antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

Anthrax Edema Factor Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anthrax Edema Factor Antibody - Protein Information**Name** cya**Function**

Edema factor (EF), which constitutes one of the three proteins composing the anthrax toxin, causes edema in the host (PubMed: [6285339](http://www.uniprot.org/citations/6285339) target="_blank">6285339, PubMed: [2108958](http://www.uniprot.org/citations/2108958) target="_blank">2108958, PubMed: [11807546](http://www.uniprot.org/citations/11807546) target="_blank">11807546). Acts as a calmodulin-dependent adenyl cyclase by converting ATP to cAMP, leading to dramatic elevation of intracellular cAMP levels in the host, thereby causing edema (PubMed: [6285339](http://www.uniprot.org/citations/6285339) target="_blank">6285339)

target="_blank">6285339, PubMed:2108958, PubMed:11807546). EF is not toxic by itself and only acts as an edema factor when associated with protective antigen (PA) to form the edema toxin (EdTx) (PubMed:11553601, PubMed:2108958). Required for the survival of germinated spores within macrophages at the early stages of infection (PubMed:11737637).

Cellular Location

Secreted. Host cytoplasm, host cytosol Note=Translocation into host cytosol is mediated via interaction with the cleaved form of protective antigen (PA-63): following secretion, EF binds via its N-terminal region to the upper rim of the ring-shaped homooligomer (homoheptamer or homooctamer) formed by PA-63 on the host cell membrane (PubMed:32047164). In this PA-63 pre-pore state, the N- terminal segment of EF refolds into an alpha helix engaged in the alpha-clamp of the PA-63 pre-pore (PubMed:32047164, PubMed:32521227) Recruitment to the PA-63 pre-pore enables domain reorganization of EF (PubMed:32521227). Loaded complexes are then endocytosed, followed by a conformational change of oligomerized PA-63 from the pre-pore to pore state, which is triggered by the low pH in the endosome (PubMed:11207582). EF is then unfolded to pass through the PA-63 pore and translocate into the host cytosol (PubMed:32047164)

Anthrax Edema Factor Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anthrax Edema Factor Antibody - Images

Anthrax Edema Factor Antibody - Background

Anthrax Edema Factor Antibody: Anthrax infection is initiated by the inhalation, ingestion, or cutaneous contact with *Bacillus anthracis* endospores. *B. anthracis* produces three polypeptides that comprise the anthrax toxin: protective antigen (PA), lethal factor (LF), and edema factor (EF). PA binds to two related proteins on the cell surface; these are termed tumor epithelial marker 8 (TEM8)/anthrax toxin receptor (ATR) and capillary morphogenesis protein 2 (CMG2), although it is still unclear which is physiologically relevant. Following PA binding to its receptor, PA is cleaved into two fragments by a furin-like protease. The bound fragment binds both LF and EF; the resulting complex is then endocytosed which allows the translocation of LF and EF into the cytoplasm. EF is a calmodulin and Ca⁺⁺-dependent adenylate cyclase responsible for the edema seen in the disease. It is thought to benefit the *B. anthracis* bacteria by inhibiting cells of the host immune system.

Anthrax Edema Factor Antibody - References

Schwartz MN. Recognition and management of anthrax - an update. *New Engl. J. Med.* 2001; 345:1621-6.
Moayeri M and Leppla SH. The roles of anthrax toxin in pathogenesis. *Curr. Opin. Microbiol.* 2004; 7:19-24.
Bradley KA, Mogridge J, Mourez M, et al. Identification of the cellular receptor for anthrax toxin.

Nature 2001; 414:225-9.

Scobie HM, Rainey GJ, Bradley KA, et al. Human capillary morphogenesis protein 2 functions as an anthrax toxin receptor. Proc. Natl. Acad. Sci. USA 2003; 100:5170-4.