

## **CTRP5 Antibody**

Catalog # ASC10340

# Specification

# **CTRP5 Antibody - Product Information**

Application WB
Primary Accession Q9BXIO

Other Accession <u>AAQ88749</u>, <u>37181891</u>

Reactivity
Host
Clonality
Polyclonal
Isotype
Human
Rabbit
Polyclonal

Application Notes CTRP5 antibody can be used for the

detection of CTRP5 by Western blot at 1 - 4

μg/mL.

# **CTRP5 Antibody - Additional Information**

Gene ID 114902

**Other Names** 

CTRP5 Antibody: CTRP5, CTRP5, UNQ303/PRO344, Complement C1q tumor necrosis factor-related protein 5, C1q and tumor necrosis factor related protein 5

#### Target/Specificity

C1QTNF5; Adipokine proteins are often highly modified post-translationally and migrate in SDS-PAGE at positions other than their predicted size.

### **Reconstitution & Storage**

CTRP5 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

#### **Precautions**

CTRP5 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## **CTRP5 Antibody - Protein Information**

Name C1QTNF5

**Synonyms** CTRP5

**Cellular Location** 

Secreted.


#### **CTRP5 Antibody - Protocols**



Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

#### CTRP5 Antibody - Images



Western blot analysis of CTRP5 in caco-2 cell lysate with CTRP5 antibody at (A) 1, (B) 2, and (C) 4  $\mu$ g/mL.

#### CTRP5 Antibody - Background

CTRP5 Antibody: Adipose tissue of an organism plays a major role in regulating physiologic and pathologic processes such as metabolism and immunity by producing and secreting a variety of bioactive molecules termed adipokines. One highly conserved family of adipokines is adiponectin/ACRP30 and its structural and functional paralogs, the C1q/tumor necrosis factor-alpha-related proteins (CTRPs) 1-7. Unlike adiponectin, which is expressed exclusively by differentiated adipocytes, the CTRPs are expressed in a wide variety of tissues. These proteins are thought to act mainly on liver and muscle tissue to control glucose and lipid metabolism. An analysis of the crystal structure of adiponectin revealed a structural and evolutionary link between TNF and C1q-containing proteins, suggesting that these proteins arose from a common ancestral innate immunity gene. CTRP5 has been suggested to be involved in age-related macular degeneration.

## **CTRP5 Antibody - References**

CTRP5 antibody can be stored at 4°C, stable for one year. Fantuzzi G. Adipose tissue, adipokines, and inflammation. J. Allergy Clin. Immunol. 2005; 115:911-9.

Tsao T-S, Lodish HF, and Fruebis J. ACRP30, a new hormone controlling fat and glucose metabolism. Euro. J. Pharmacol. 2002; 440:213-21.

Wong GW, Wang J, Hug C, et al. A family of Acrp30/ adiponectin structural and functional paralogs. Proc. Natl. Acad. Sci. USA 2004; 101:10302-7.

Shapiro L and Scherer PE. The crystal structure of a complement-1q family protein suggests an evolutionary link to tumor necrosis factor. Curr. Biol. 1998; 8:335-8.