

**RP105 Antibody**  
Catalog # ASC10429**Specification**

---

**RP105 Antibody - Product Information**

Application	WB, IHC, IF
Primary Accession	<a href="#">Q99467</a>
Other Accession	<a href="#">BAA12019</a> , <a href="#">1843411</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	RP105 antibody can be used for the detection of RP105 by Western blot at 0.5 - 1 µg/mL. Despite its predicted molecular weight, RP105 often migrates at 95 - 105 kDa. Antibody can also be used for immunohistochemistry starting at 2 µg/mL. For immunofluorescence start at 20 µg/mL.

**RP105 Antibody - Additional Information**

Gene ID	4064
<b>Other Names</b>	
RP105 Antibody: LY64, Ly78, RP105, LY64, CD180 antigen, Lymphocyte antigen 64, CD180 molecule	

**Target/Specificity**  
CD180;**Reconstitution & Storage**

RP105 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

**Precautions**

RP105 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**RP105 Antibody - Protein Information****Name** CD180**Synonyms** LY64, RP105**Function**

May cooperate with MD-1 and TLR4 to mediate the innate immune response to bacterial lipopolysaccharide (LPS) in B-cells. Leads to NF- kappa-B activation. Also involved in the life/death decision of B-cells (By similarity).

### Cellular Location

Cell membrane; Single-pass type I membrane protein

### Tissue Location


Expressed mainly on mature peripheral B cells. Detected in spleen, lymph node and appendix.  
Not detected in pre-B and -T cells

### RP105 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### RP105 Antibody - Images



Western blot analysis of RP105 in human spleen tissue lysate with RP105 antibody at (A) 0.5 and (B) 1 µg/mL.



Immunohistochemistry of RP105 in human spleen tissue with RP105 antibody at 2 µg/mL.



Immunofluorescence of RP105 in human spleen tissue with RP105 antibody at 20 µg/mL.

### **RP105 Antibody - Background**

RP105 Antibody: Toll-like receptors (TLRs) are evolutionarily conserved pattern-recognition molecules resembling the toll proteins that mediate antimicrobial responses in *Drosophila*. These proteins recognize different microbial products during infection and serve as an important link between the innate and adaptive immune responses. The signaling of these TLRs is kept under tight control by the expression of endogenous inhibiting proteins. One such protein is RP105, a recently identified homolog to TLR4 that, with MD-1, interacts with and inhibits the TLR4/MD-2 signaling pathway. It has also been suggested that the RP105/MD-1 complex influences antibody production mediated by both TLR4/MD-2 and TLR2 receptor complexes.

### **RP105 Antibody - References**

- Takeda K, Kaisho T, and Akira S. Toll-like receptors. *Annu. Rev. Immunol.* 2003; 21:335-76.  
Janeway CA Jr. and Medzhitov R. Innate immune recognition. *Annu. Rev. Immunol.* 2002; 20:197-216.  
Divanovic S, Trompette A, Atabani SF, et al. Inhibition of TLR-4/MD-2 signaling by RP105/MD-1. *J. Endotoxin Res.* 2005; 11:363-8.  
Nagai Y, Kobayashi T, Motoi Y, et al. The radioprotective 105/MD-1 complex links TLR2 and TLR4/MD-2 in antibody response to microbial membranes. *J. Immunol.* 2005; 174:7043-9.