

TRPC6 Antibody

Catalog # ASC10451

Specification

TRPC6 Antibody - Product Information

Application
Primary Accession
Other Accession
Reactivity
Host
Clonality
Isotype
Application Notes

WB, IHC-P, IF, E
O9Y210
AAH93658, 7225
Human, Mouse
Rabbit
Polyclonal

TRPC6 antibody can be used for the detection of TRPC6 by Western blot at 0.5 - 1 μ g/mL. Antibody can also be used for Immunohistochemistry starting at 10 μ g/mL and Immunocytochemistry starting at 2 μ g/mL. For immunofluorescence start at 5 μ g/mL.

TRPC6 Antibody - Additional Information

Gene ID **7225**

Other Names

TRPC6 Antibody: TRP6, FSGS2, TRP6, Short transient receptor potential channel 6, Transient receptor protein 6, TrpC6, transient receptor potential cation channel, subfamily C, member 6

Target/Specificity

TRPC6 antibody was raised against a 14 amino acid synthetic peptide from near the amino terminus of human TRPC6.

The immunogen is located within amino acids 50 - 100 of TRPC6.

Reconstitution & Storage

TRPC6 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

TRPC6 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.


TRPC6 Antibody - Protein Information

Name TRPC6 {ECO:0000303|PubMed:9930701, ECO:0000312|HGNC:HGNC:12338}

Function

Forms a receptor-activated non-selective calcium permeant cation channel (PubMed:19936226, PubMed:23291369, PubMed:<a hre







href="http://www.uniprot.org/citations/26892346" target=" blank">26892346, PubMed:9930701). Probably is operated by a phosphatidylinositol second messenger system activated by receptor tyrosine kinases or G-protein coupled receptors. Activated by diacylglycerol (DAG) in a membranedelimited fashion, independently of protein kinase C (PubMed: 26892346). Seems not to be activated by intracellular calcium store depletion.

Cellular Location

Cell membrane; Multi-pass membrane protein

Tissue Location


Expressed primarily in placenta, lung, spleen, ovary and small intestine. Expressed in podocytes and is a component of the glomerular slit diaphragm.

TRPC6 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

TRPC6 Antibody - Images



Immunohistochemistry of CHADL in mouse skeletal muscle tissue with CHADL antibody at 5 μg/mL.





TRPC6 Antibody - Background

TRPC6 Antibody: The mammalian transient receptor potential (TRP) superfamily can be divided into three major families including the "canonical TRP" (TRPC) family. The seven members of this family share the activation through PLC-coupled receptors and have been suggested to be components of receptor-regulated cation channels in different cell types. Furthermore, the members of the TRPC3/6/7 subfamily can be activated by diacylglycerol (DAG) analogs, suggesting a possible mechanism of activation of these channels by PLC-coupled receptors. When expressed in transfected cells, TRPC6 acts as a non-selective store-independent receptor-activated cation channel. It is activated by DAG in a PKC-independent manner and is insensitive to IP3 activation. There is increasing evidence that TRPC6 encodes endogenous DAG-activated receptor-operated cation channels in vivo.

TRPC6 Antibody - References

Contell C, Birnbaumer V, Flockerzi V, et al. A unified nomenclature for the superfamily of TRP cation channels. Mol. Cell 2002; 9:229-31.

Dietrich A, Kalwa H, Rost BR, et al. The diacylglycerol-sensitive TRPC3/6/7 subfamily of cation channels: functional characterization and physiological relevance. Pflugers Arch. 2005; 451:72-80. Trebak M, Vazquez G, Bird GSJ, et al. The TRPC3/6/7 subfamily of cation channels. Cell Calcium 2003; 33:451-61