

**ORAI1 Antibody**  
**Catalog # ASC10502****Specification**

---

**ORAI1 Antibody - Product Information**

Application	WB, IHC-P, IF, E
Primary Accession	<a href="#">Q96D31</a>
Other Accession	<a href="#">Q96D31</a> , <a href="#">84876</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	Predicted: 33 kDa

Application Notes	<b>Observed: 50 kDa KDa</b> <b>ORAI1 antibody can be used for detection of ORAI1 by Western blot at 1 µg/mL. Antibody can also be used for immunohistochemistry starting at 10 µg/mL. For immunofluorescence start at 20 µg/mL.</b>
-------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

**ORAI1 Antibody - Additional Information**Gene ID **84876****Other Names**

ORAI1 Antibody: IMD9, ORAT1, CRACM1, TMEM142A, Calcium release-activated calcium channel protein 1, Protein orai-1, ORAI calcium release-activated calcium modulator 1

**Target/Specificity**

ORAI1 antibody was raised against an 18 amino acid synthetic peptide from near the amino terminus of human ORAI1.<br><br>The immunogen is located within the first 50 amino acids of ORAI1.

**Reconstitution & Storage**

ORAI1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

**Precautions**

ORAI1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**ORAI1 Antibody - Protein Information****Name** ORAI1 {ECO:0000303|PubMed:16921383, ECO:0000312|HGNC:HGNC:25896}**Function**

Pore-forming subunit of two major inward rectifying Ca(2+) channels at the plasma membrane:

Ca(2+) release-activated Ca(2+) (CRAC) channels and arachidonate-regulated Ca(2+)-selective (ARC) channels (Probable) (PubMed:<a href="http://www.uniprot.org/citations/16645049" target="\_blank">16645049</a>, PubMed:<a href="http://www.uniprot.org/citations/16733527" target="\_blank">16733527</a>, PubMed:<a href="http://www.uniprot.org/citations/16807233" target="\_blank">16807233</a>, PubMed:<a href="http://www.uniprot.org/citations/16921383" target="\_blank">16921383</a>, PubMed:<a href="http://www.uniprot.org/citations/19249086" target="\_blank">19249086</a>, PubMed:<a href="http://www.uniprot.org/citations/19706554" target="\_blank">19706554</a>, PubMed:<a href="http://www.uniprot.org/citations/23307288" target="\_blank">23307288</a>, PubMed:<a href="http://www.uniprot.org/citations/26956484" target="\_blank">26956484</a>, PubMed:<a href="http://www.uniprot.org/citations/28219928" target="\_blank">28219928</a>). Assembles with ORAI2 and ORAI3 to form hexameric CRAC channels that mediate Ca(2+) influx upon depletion of endoplasmic reticulum Ca(2+) store and channel activation by Ca(2+) sensor STIM1, a process known as store-operated Ca(2+) entry (SOCE). Various pore subunit combinations may account for distinct CRAC channel spatiotemporal and cell-type specific dynamics. ORAI1 mainly contributes to the generation of Ca(2+) plateaus involved in sustained Ca(2+) entry and is dispensable for cytosolic Ca(2+) oscillations, whereas ORAI2 and ORAI3 generate oscillatory patterns. CRAC channels assemble in Ca(2+) signaling microdomains where Ca(2+) influx is coupled to calmodulin and calcineurin signaling and activation of NFAT transcription factors recruited to ORAI1 via AKAP5. Activates NFATC2/NFAT1 and NFATC3/NFAT4-mediated transcriptional responses. CRAC channels are the main pathway for Ca(2+) influx in T cells and promote the immune response to pathogens by activating NFAT-dependent cytokine and chemokine transcription (PubMed:<a href="http://www.uniprot.org/citations/16582901" target="\_blank">16582901</a>, PubMed:<a href="http://www.uniprot.org/citations/17442569" target="\_blank">17442569</a>, PubMed:<a href="http://www.uniprot.org/citations/19182790" target="\_blank">19182790</a>, PubMed:<a href="http://www.uniprot.org/citations/20354224" target="\_blank">20354224</a>, PubMed:<a href="http://www.uniprot.org/citations/22641696" target="\_blank">22641696</a>, PubMed:<a href="http://www.uniprot.org/citations/26221052" target="\_blank">26221052</a>, PubMed:<a href="http://www.uniprot.org/citations/32415068" target="\_blank">32415068</a>, PubMed:<a href="http://www.uniprot.org/citations/33941685" target="\_blank">33941685</a>). Assembles with ORAI3 to form channels that mediate store-independent Ca(2+) influx in response to inflammatory metabolites arachidonate or its derivative leukotriene C4, termed ARC and LRC channels respectively (PubMed:<a href="http://www.uniprot.org/citations/19622606" target="\_blank">19622606</a>, PubMed:<a href="http://www.uniprot.org/citations/32415068" target="\_blank">32415068</a>). Plays a prominent role in Ca(2+) influx at the basolateral membrane of mammary epithelial cells independently of the Ca(2+) content of endoplasmic reticulum or Golgi stores. May mediate transepithelial transport of large quantities of Ca(2+) for milk secretion (By similarity) (PubMed:<a href="http://www.uniprot.org/citations/20887894" target="\_blank">20887894</a>).

### Cellular Location

Cell membrane; Multi-pass membrane protein. Basolateral cell membrane {ECO:0000250|UniProtKB:Q8BWG9}; Multi-pass membrane protein. Note=Upon store depletion, colocalizes with STIM1 in membrane punctae at ER-PM junctions (PubMed:19182790, PubMed:19249086, PubMed:26221052, PubMed:27185316) [Isoform beta]; Cell membrane

### Tissue Location

Expressed in naive CD4 and CD8 T cells (at protein level) (PubMed:26956484). Expressed at similar levels in naive and effector T helper cells (PubMed:20354224)


## ORAI1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### ORAI1 Antibody - Images



Western blot analysis of TEM4 in Jurkat cell lysate with TEM4 antibody at (A) 0.5 and (B) 1  $\mu$ g/mL.

### ORAI1 Antibody - Background

ORAI1 Antibody: Antigen stimulation of immune cells triggers  $\text{Ca}^{++}$  entry through  $\text{Ca}^{++}$  release-activated  $\text{Ca}^{++}$  (CRAC) channels. ORAI1 is a recently identified four-transmembrane spanning protein that is an essential component of CRAC. A missense mutation in this protein in humans is the cause of one form of hereditary severe combined immune deficiency (SCID) which results in ablated T-cell  $\text{Ca}^{++}$  entry. It has been suggested that ORAI1 functions as a highly selective  $\text{Ca}^{++}$  plasma membrane channel that is gated through interactions with STIM1, the store-activated endoplasmic reticulum  $\text{Ca}^{++}$  sensor.

### ORAI1 Antibody - References

- Lewis RS. Calcium signaling mechanisms in T lymphocytes. *Annu. Rev. Immunol.* 2001; 19:497-521.
- Feske S, Gwack Y, Prakriya M, et al. A mutation in Orai1 causes immune deficiency by abrogating CRAC channel function. *Nature* 2006; 441:179-85.
- Soboloff J, Spassova MA, Dziadek MA, et al. Calcium signals mediated by STIM and Orai proteins - a new paradigm in inter-organelle communication. *Biochim. Biophys. Acta.* 2006; 1763:1161-8.