

**PD-1 Antibody**  
**Catalog # ASC10512****Specification**

---

**PD-1 Antibody - Product Information**

Application	WB, IHC-P, E
Primary Accession	<a href="#">Q15116</a>
Other Accession	<a href="#">Q15116</a> , <a href="#">145559515</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	Predicted: 32 kDa

Application Notes	<b>Observed: 43 kDa</b> PD-1 antibody can be used for detection of PD-1 by Western blot at 1 µg/mL. Antibody can also be used for immunohistochemistry starting at 5 µg/mL.
-------------------	--

**PD-1 Antibody - Additional Information**

Gene ID **5133**

**Other Names**

PD-1 Antibody: PD1, PD-1, CD279, SLEB2, hPD-1, hPD-I, hSLE1, PD1, Programmed cell death protein 1, Protein PD-1, programmed cell death 1

**Target/Specificity**

PDCD1;

**Reconstitution & Storage**

PD-1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

**Precautions**

PD-1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**PD-1 Antibody - Protein Information**

**Name** PDCD1 {ECO:0000303|PubMed:7851902, ECO:0000312|HGNC:HGNC:8760}

**Function**

Inhibitory receptor on antigen activated T-cells that plays a critical role in induction and maintenance of immune tolerance to self (PubMed:<a href="http://www.uniprot.org/citations/21276005" target="\_blank">21276005</a>, PubMed:<a href="http://www.uniprot.org/citations/37208329" target="\_blank">37208329</a>). Delivers

inhibitory signals upon binding to ligands CD274/PDCD1L1 and CD273/PDCD1LG2 (PubMed:<a href="http://www.uniprot.org/citations/21276005" target="\_blank">21276005</a>). Following T-cell receptor (TCR) engagement, PDCD1 associates with CD3- TCR in the immunological synapse and directly inhibits T-cell activation (By similarity). Suppresses T-cell activation through the recruitment of PTPN11/SHP-2: following ligand-binding, PDCD1 is phosphorylated within the ITSM motif, leading to the recruitment of the protein tyrosine phosphatase PTPN11/SHP-2 that mediates dephosphorylation of key TCR proximal signaling molecules, such as ZAP70, PRKCQ/PKCtheta and CD247/CD3zeta (By similarity).

#### Cellular Location


Cell membrane; Single-pass type I membrane protein

#### PD-1 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### PD-1 Antibody - Images



Western blot analysis of PD-1 in THP-1 cell lysate with PD-1 antibody at 1 µg/mL in the (A) absence and (B) presence of blocking peptide.



Immunohistochemistry of PD-1 in human brain tissue with PD-1 antibody at 5 µg/mL.

#### PD-1 Antibody - Background

**PD-1 Antibody:** Cell-mediated immune responses are initiated by T lymphocytes that are themselves stimulated by cognate peptides bound to MHC molecules on antigen-presenting cells (APC). T-cell activation is generally self-limited as activated T cells express receptors such as PD-1 (also known as PDCD-1) that mediate inhibitory signals from the APC. PD-1 can bind two different but related ligands, PDL-1 and PDL-2. Upon binding to either of these ligands, signals generated by PD-1 inhibit the activation of the immune response in the absence of "danger signals" such as LPS or other molecules associated with bacteria or other pathogens. Evidence for this is seen in PD1-null mice who exhibit hyperactivated immune systems and autoimmune diseases.

#### **PD-1 Antibody - References**

Holling TM, Schooten E, and van Den Elsing PJ. Function and regulation of MHC class II molecules in T-lymphocytes: of mice and men. *Hum. Immunol.* 2004; 65:282-90.

Ishida Y, Agata Y, Shibahara K, et al. Induced expression of PD-1, a novel member of the immunoglobulin gene superfamily, upon programmed cell death. *EMBO J.* 1992; 11:3887-95.

Zhong X, Bai C, Gao W, et al. Suppression of expression and function of negative immune regulator PD-1 by certain pattern recognition and cytokine receptor signals associated with immune system danger. *Int. Immunol.* 2004; 16:1181-8.

Nishimura H, Nose M, Hiai H, et al. Development of lupus-like autoimmune diseases by the disruption of the PD-1 gene encoding an ITIM motif-carrying immunoreceptor. *Immunity* 1999; 11:141-51.