

AIFM3 Antibody
Catalog # ASC10668**Specification**

AIFM3 Antibody - Product Information

Application	WB
Primary Accession	Q96NN9
Other Accession	Q96NN9 , 74732608
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	AIFM3 antibody can be used for the detection of AIFM3 by Western blot at 1 - 2 µg/mL.

AIFM3 Antibody - Additional Information

Gene ID	150209
Target/Specificity	
AIFM3;	

Reconstitution & Storage

AIFM3 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

AIFM3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

AIFM3 Antibody - Protein Information

Name AIFM3

Synonyms AIFL

Function

Induces apoptosis through a caspase dependent pathway. Reduces mitochondrial membrane potential.

Cellular Location

Mitochondrion. Note=Does not translocate to the nucleus upon induction of apoptosis

Tissue Location


Ubiquitous. Expressed in bone marrow, cerebral cortex, liver, ovary, thymus, thyroid gland and tongue (at protein level).

AIFM3 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

AIFM3 Antibody - Images



Western blot analysis of AIFM3 in human brain tissue lysate with AIFM3 antibody at (A) 1 and (B) 2 µg/mL.

AIFM3 Antibody - Background

AIFM3 Antibody: Apoptosis, also known as programmed cell death, plays major roles in development and normal tissue turnover in addition to tumor formation. Recently a protein similar to the apoptosis-inducing factor (AIF) was cloned and designated AIFL (also known as AIFM3). AIFM3 is expressed ubiquitously and is predominantly localized to the inner membranes of mitochondria. Unlike AIF, AIFM3 does not translocate to the nucleus upon induction of apoptosis. However, overexpression of AIFM3, like AIF, induced cytochrome c release from the mitochondria, cleavage of caspase 3, and ultimately apoptosis, indicating AIFM3 induces apoptosis through caspase activation. Multiple isoforms of AIFM3 are known to exist.

AIFM3 Antibody - References

Jin Z and El Deiry WS. Overview of cell death signaling pathways. *Cancer Biol. Ther.*2004; 4:139-63
Xie Q, Lin T, Zhang Y, et al. Molecular cloning and characterization of a human AIF-like gene with the ability to induce apoptosis. *J. Biol. Chem.*2005; 280:19673-81.