

## **RGPD5 Antibody**

Catalog # ASC10684

# **Specification**

# **RGPD5 Antibody - Product Information**

Application WB
Primary Accession Q99666
Other Accession NP 005045, 83267877

Reactivity
Host
Clonality
Human
Rabbit
Polyclonal

Isotype

Application Notes

RGPD5 antibody can be used for detection of RGPD5 by Western blot at 1 µg/mL.

## **RGPD5 Antibody - Additional Information**

Gene ID **84220** 

Target/Specificity

RGPD5;

#### **Reconstitution & Storage**

RGPD5 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

#### **Precautions**

RGPD5 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

### **RGPD5 Antibody - Protein Information**

Name RGPD5

Synonyms RANBP2L1, RGP5, RGP7, RGPD7

**Cellular Location** 

Cytoplasm.

**Tissue Location** 

Expressed in testis..

#### **RGPD5 Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

Western Blot



- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# **RGPD5 Antibody - Images**



Western blot analysis of RGPD5 in human thymus tissue lysate with RGPD5 antibody at 1  $\mu$ g/mL in (A) the absence and (B) the presense of blocking peptide.

# **RGPD5 Antibody - Background**

RGPD5 Antibody: The RANBP2-like and GRIP domain containing 5 protein (RGPD5) has high similarity to RANBP2, a large RAN-binding protein localized at the cytoplasmic side of the nuclear pore complex. The gene coding for RGPD5 is thought to have arisen from a gene duplication event of RANBP2 as these highly homologous genes are located close to each other at chromosome 2q11-q12. RGPD5 was identified as an HIV dependency factor (HDF), suggesting that RGPD5 may be an important drug target in HIV treatment. At least two isoforms of RGPD5 are known to exist, of which the shorter isoform is expressed primarily in testis, while the longer of the two is expressed at low levels in a number of somatic tissues.

### **RGPD5 Antibody - References**

Nothwang HG, Rensing C, Kubler M, et al. Identification of a novel Ran binding protein 2 related gene (RANBP2L1) and detection of a gene cluster on chromosome 2q11-q12. Genomics1998; 47:383-92.

Brass AL, Dykxhoorn DM, Benita Y, et al. Identification of host proteins required for HIV infection through a functional genomic screen. Science2008; 319:921-6.

Wang LF, Zhu HD, Miao SY, et al. Molecular cloning and characterization of a novel testis-specific nucleoporin-related gene. Arch. Androl.1999; 42:71-84.