

NUP107 Antibody

Catalog # ASC10729

Specification

NUP107 Antibody - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Isotype Application Notes WB, IF, ICC, E P57740 P57740, 12230339 Human, Mouse, Rat Rabbit Polyclonal IgG NUP107 antibody can be used for detection of NUP107 by Western blot at 1 - 2 μg/mL. Antibody can also be used for immunocytochemistry starting at 10 μg/mL. For immunofluorescence start at 10 μg/mL.

NUP107 Antibody - Additional Information

Gene ID Target/Specificity NUP107;

Reconstitution & Storage

NUP107 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

57122

Precautions

NUP107 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

NUP107 Antibody - Protein Information

Name NUP107

Function

Plays a role in the nuclear pore complex (NPC) assembly and/or maintenance (PubMed:12552102, PubMed:15229283, PubMed:30179222). Required for the assembly of peripheral proteins into the NPC (PubMed:12552102, PubMed:12552102, PubMed:12552102, PubMed:12552102, PubMed:15229283, PubMed:15229283, PubMed:15229283, PubMed:15229283, PubMed:30179222, PubMed:30179222, PubMed:30179222, PubMed:30179222, PubMed:<a href="http://www.uniprot.or



Cellular Location

Nucleus membrane. Nucleus, nuclear pore complex. Chromosome, centromere, kinetochore. Note=Located on both the cytoplasmic and nuclear sides of the NPC core structure (PubMed:11564755). During mitosis, localizes to the kinetochores (PubMed:11564755). Dissociates from the dissasembled NPC structure late during prophase of mitosis (PubMed:11564755)

Tissue Location

Ubiquitously expressed in fetal and adult tissues.

NUP107 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

NUP107 Antibody - Images



Western blot analysis of NUP107 in A549 cell lysate with NUP107 antibody at (A) 1 and (B) 2 $\mu g/mL$



Immunocytochemistry of NUP107 in A549 cells with NUP107 antibody at 10 µg/mL.





Immunofluorescence of NUP107 in A549 cells with NUP107 antibody at 10 µg/mL.

NUP107 Antibody - Background

NUP107 Antibody: The nuclear pore complex (NPC) is a protein assembly localized at the nuclear rim and mediates macromolecular transport between the nucleus and the cytoplasm. The mammalian nucleoporin (NUP)-107 is part of the hetero-oligomeric complex that also contains NUP160, NUP133, NUP96, and mammalian homolog of yeast sec13p. While the majority of the NUP107-160 nuclear pore sub-complex localizes to the nuclear pore, a small fraction is observed at kinetochores and pro-metaphase spindle poles in mitotic cells in association with proteins such as Mad1, Mad2, Bub3 and Cdc20. Immunodepletion of the NUP107-160 complex resulted in defective spindle assembly indicating that it has multiple functions. NUP107 has recently been identified as an HIV dependency factor (HDF), suggesting that NUP107 may be an important drug target in HIV treatment. Multiple isoforms of NUP107 are known to exist.

NUP107 Antibody - References

Tran EJ and Wente SR. Dynamic nuclear pore complex: life on the edge. Cell2006; 125:1041-53. Boehmer T, Enninga J, Dales S, et al. Depletion of a single nucleoporin, Nup107, prevents the assembly of a subset of nucleoporins into the nuclear pore complex. Proc. Natl. Acad. Sci. USA2003; 100:981-5.

Orjalo AV, Arnaoutov A, Shen Z, et al. The Nup107-160 nucleoporin complex is required for correct bipolar spindle assembly. Mol. Bio. Cell2006; 17:3806-18.

Brass AL, Dykxhoorn DM, Benita Y, et al. Identification of host proteins required for HIV infection through a functional genomic screen. Science2008; 319:921-6.