

TREX1 Antibody
Catalog # ASC10826**Specification**

TREX1 Antibody - Product Information

Application	WB, IHC-P, IF, E
Primary Accession	Q9NSU2
Other Accession	NP_057465 , 7705353
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	Predicted: 33, 35, 41 kDa

Application Notes	Observed: 31 kDa KDa TREX1 antibody can be used for detection of TREX1 by Western blot at 0.5 and 1 µg/mL. Antibody can also be used for immunohistochemistry starting at 2.5 µg/mL. For immunofluorescence start at 20 µg/mL.
-------------------	---

TREX1 Antibody - Additional InformationGene ID **11277****Target/Specificity**

TREX1; TREX1 antibody will not cross-react with the related protein TREX2. At least three isoforms of TREX1 are known to exist; this antibody will recognize all three isoforms.

Reconstitution & Storage

TREX1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

TREX1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

TREX1 Antibody - Protein Information

Name TREX1 {ECO:0000303|PubMed:10391904, ECO:0000312|HGNC:HGNC:12269}

Function

Major cellular 3'-to-5' DNA exonuclease which digests single- stranded DNA (ssDNA) and double-stranded DNA (dsDNA) with mismatched 3' termini (PubMed:10391904, PubMed:10393201, PubMed:17293595). Prevents cell-intrinsic initiation of autoimmunity (PubMed:17293595).

href="http://www.uniprot.org/citations/10391904" target="_blank">10391904, PubMed:10393201, PubMed:17293595). Acts by metabolizing DNA fragments from endogenous retroelements, including L1, LTR and SINE elements (PubMed:10391904, PubMed:10393201, PubMed:17293595). Plays a key role in degradation of DNA fragments at cytosolic micronuclei arising from genome instability: its association with the endoplasmic reticulum membrane directs TREX1 to ruptured micronuclei, leading to micronuclear DNA degradation (PubMed:33476576). Micronuclear DNA degradation is required to limit CGAS activation and subsequent inflammation (PubMed:33476576). Unless degraded, these DNA fragments accumulate in the cytosol and activate the cGAS-STING innate immune signaling, leading to the production of type I interferon (PubMed:33476576). Prevents chronic ATM-dependent checkpoint activation, by processing ssDNA polynucleotide species arising from the processing of aberrant DNA replication intermediates (PubMed:18045533). Inefficiently degrades oxidized DNA, such as that generated upon antimicrobial reactive oxygen production or upon absorption of UV light (PubMed:23993650). During GZMA-mediated cell death, contributes to DNA damage in concert with NME1 (PubMed:16818237). NME1 nicks one strand of DNA and TREX1 removes bases from the free 3' end to enhance DNA damage and prevent DNA end reannealing and rapid repair (PubMed:16818237).

Cellular Location

Nucleus. Cytoplasm, cytosol. Endoplasmic reticulum membrane; Peripheral membrane protein. Note=Retained in the cytoplasm through the C-terminal region (By similarity). Localization to the endoplasmic reticulum membrane is required to direct TREX1 to ruptured micronuclei (PubMed:33476576). In response to DNA damage, translocates to the nucleus where it is specifically recruited to replication foci (PubMed:16818237). Translocation to the nucleus also occurs during GZMA-mediated cell death (PubMed:16818237) {ECO:0000250|UniProtKB:Q91XB0, ECO:0000269|PubMed:16818237, ECO:0000269|PubMed:33476576}

Tissue Location

Detected in thymus, spleen, liver, brain, heart, small intestine and colon.


TREX1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TREX1 Antibody - Images






Western blot analysis of TREX1 in human spleen tissue lysate with TREX1 antibody at (A) 0.5 and (B) 1 μ g/mL.



Immunohistochemistry of TREX1 in human spleen tissue with TREX1 antibody at 2.5 μ g/mL.



Immunofluorescence of TREX1 in Human Spleen tissue with TREX1 antibody at 20 μ g/mL.

TREX1 Antibody - Background

TREX1 Antibody: Trex1 is the major human 3' to 5' exonuclease which is required for checkpoint signaling after DNA damage. It is ubiquitously expressed, binds to single stranded DNA coated with replication protein A that accumulates at sites of DNA damage and recruits the ataxia telangiectasia and Rad3 related protein (ATR), a checkpoint kinase, to sites of DNA damage and replication stress. Trex1 is required for ATR expression. This gene uses two different open reading frames. The upstream ORF encodes proteins which interact with ATR and localize to intranuclear foci induced by DNA damage and are essential components of the DNA damage checkpoint. The downstream ORF encodes proteins with 3' to 5' exonuclease activity and may be a subunit of human DNA polymerase III. Mutations in this gene result in Aicardi-Goutieres syndrome, chilblain

lupus, and Cree encephalitis.

TREX1 Antibody - References

Mazur DJ and Perrino FW. Identification and expression of the TREX1 and TREX2 cDNA sequences encoding mammalian 3→5 exonucleases. J. Biol. Chem. 1999; 274:19655-60.

Cortez D, Guntuku S, Qin J, et. al. ATR and ATRIP: partners in checkpoint signaling. Science 2001 294:1713-6

Stetson DB, Ko JS, Heidmann T, et al. Trex1 prevents cell-intrinsic initiation of autoimmunity. Cell 2008 134:587-98.

Mazur DJ and Perrino FW. Structure and expression of the TREX1 and TREX2 3'→5' exonuclease genes. J. Biol. Chem. 2001; 276:14718-27.