

LIAR Antibody
Catalog # ASC10942**Specification**

LIAR Antibody - Product Information

Application	WB, IHC-P, IF, E
Primary Accession	Q6NXT1
Other Accession	NP_620152 , 20270347
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	LIAR antibody can be used for detection of LIAR by Western blot at 1 µg/mL. Antibody can also be used for immunohistochemistry starting at 2.5 µg/mL. For immunofluorescence start at 20 µg/mL.

LIAR Antibody - Additional Information

Gene ID	129138
Target/Specificity	
ANKRD54;	

Reconstitution & Storage

LIAR antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

LIAR Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

LIAR Antibody - Protein Information

Name ANKRD54

Synonyms LIAR

Function

Plays an important role in regulating intracellular signaling events associated with erythroid terminal differentiation.

Cellular Location


Nucleus. Cytoplasm. Midbody. Note=Shuttles between nucleus and cytoplasm during the cell cycle. EPO stimulation induces nuclear accumulation (By similarity).

LIAR Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


LIAR Antibody - Images



Western blot analysis of LIAR in mouse kidney tissue lysate with LIAR antibody at 1 μ g/mL in (A) the absence and (B) the presence of blocking peptide.



Immunohistochemistry of LIAR in mouse kidney tissue with LIAR antibody at 2.5 μ g/mL.



Immunofluorescence of LIAR in Mouse Kidney cells with LIAR antibody at 20 μ g/mL.

LIAR Antibody - Background

LIAR Antibody: Ankyrin (ANK) repeats mediate protein-protein interactions in diverse families of proteins. The number of ANK repeats in a protein can range from 2 to over 20. ANK repeats may occur in combinations with other types of domains. The structural repeat unit contains two anti-parallel helices and a beta-hairpin, with repeats stacked in a superhelical arrangement. LIAR, also known as ANKRD54, is a recently identified ANK repeat-containing protein that is predominantly expressed in tissues rich in ciliated cells, such as olfactory sensory neurons and is predicted to be important to cilia. At least three isoforms of LIAR are known to exist.

LIAR Antibody - References

Li J, Mahajan A, and Tsai MD. Ankyrin repeat: a unique motif mediating protein-protein interactions. *Biochemistry* 2006; 45:15168-78.