

PPAPDC3 Antibody
Catalog # ASC11029**Specification**

PPAPDC3 Antibody - Product Information

| | |
|-------------------|---|
| Application | WB, IF |
| Primary Accession | Q8NBV4 |
| Other Accession | EAW87966 , 119608372 |
| Reactivity | Human, Mouse, Rat |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | IgG |
| Application Notes | PPAPDC3 antibody can be used for detection of PPAPDC3 by Western blot at 1 - 2 µg/mL. For immunofluorescence start at 20 µg/mL. |

PPAPDC3 Antibody - Additional Information

| | |
|---------------------------|-------|
| Gene ID | 84814 |
| Target/Specificity | |
| PPAPDC3; | |

Reconstitution & Storage

PPAPDC3 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

PPAPDC3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

PPAPDC3 Antibody - Protein Information

Name PLPP7 ([HGNC:28174](#))

Synonyms C9orf67, PPAPDC3

Function

Plays a role as negative regulator of myoblast differentiation, in part through effects on MTOR signaling. Has no detectable enzymatic activity (By similarity).

Cellular Location


Nucleus envelope. Endoplasmic reticulum membrane. Membrane; Multi-pass membrane protein. Note=Both the N- and C-terminal are exposed to the cytoplasm/nucleoplasm.

PPAPDC3 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PPAPDC3 Antibody - Images



Western blot analysis of PPAPDC3 in mouse heart tissue lysate with PPAPDC3 antibody at (A) 1 and (B) 2 μ g/mL.



Immunofluorescence of PPAPDC3 in mouse heart tissue with PPAPDC3 antibody at 20 μ g/mL.

PPAPDC3 Antibody - Background

PPAPDC3 Antibody: PPAPDC3, also known as nuclear envelope transmembrane protein 39 (NET39), was initially discovered in an in silico screen for secreted or membrane proteins. It is a member of

the PAP2 superfamily of phosphatases and haloperoxidases. PPAPDC3 has recently been shown to act as a negative regulator of myoblast differentiation by diminishing the activity of the mammalian target of rapamycin TOR. PPAPDC3 is highly expressed in cardiac and skeletal muscle and becomes strongly upregulated during cultured myoblast differentiation tissues. Overexpression of PPAPDC3 in myoblasts repressed myogenesis while knockdown by RNA interference promoted differentiation indicating its part in the regulatory mechanism for myogenesis.

PPAPDC3 Antibody - References

Otsuki T, Ota T, Nishikawa T, et al. Signal sequence and keyword trap in silico for selection of full-length human cDNAs encoding secretion or membrane proteins from oligo-capped cDNA libraries. DNA Res.2005; 12:117-26.

Liu GH, Guan T, Datta K, et al. Regulation of myoblast differentiation by the nuclear envelope protein NET39. Mol. Cell Biol.2009; 29:5800-12.