

RHBDD2 Antibody

Catalog # ASC11033

Specification

RHBDD2 Antibody - Product Information

Application
Primary Accession
Other Accession
Reactivity
Host
Clonality
Isotype

Application Notes

WB, IF, E 06NTF9

NP_001035546, 94818790 Human, Mouse, Rat

Rabbit Polyclonal

IgG

RHBDD2 antibody can be used for

detection of RHBDD2 by Western blot at 1 - 2 μg/mL. For immunofluorescence start at

20 μg/mL.

RHBDD2 Antibody - Additional Information

Gene ID **57414**

Target/Specificity

RHBDD2;

Reconstitution & Storage

RHBDD2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

RHBDD2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

RHBDD2 Antibody - Protein Information

Name RHBDD2

Synonyms RHBDL7

Cellular Location

Golgi apparatus, cis-Golgi network membrane; Multi-pass membrane protein

RHBDD2 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides




- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

RHBDD2 Antibody - Images



Western blot analysis of RHBDD2 in rat lung tissue lysate with RHBDD2 antibody at (A) 1 and (B) $2 \mu g/mL$.



Immunofluorescence of RHBDD2 in Rat Lung cells with RHBDD2 antibody at 20 μg/mL.

RHBDD2 Antibody - Background

RHBDD2 Antibody: The Rhomboid family of proteins is made up of several widely conserved polytopic membrane serine proteases that play roles in growth and development. Little is known about the role of RHBDD2. Its overexpression in breast carcinoma appears to be an indicator of poor prognosis and may play a role in breast cancer progression. The related proteins RHBDD1 and RHBDD3 are both involved in the regulation of apoptosis, suggesting that RHBDD2 may also be involved in the modulation of apoptotic activity.

RHBDD2 Antibody - References

Koonin EV, Makarova KS, Rogozin IB, et al. The rhomboids: a nearly ubiquitous family of intramembrane serine proteases that probably evolved by multiple horizontal gene transfers.







Genome Biol.2003; 4:R19.

Abba MC, Sun H, Hawkins KA, et al. Breast cancer molecular signatures as determined by SAGE: correlation with lymph node status. Mol. Cancer Res.2007; 5:881-90.

Abba MC, Lacunza E, Nunez MI, et al. Rhomboid domain containing 2 (RHBDD2): a novel cancer-related gene over-expressed in breast cancer. Biochim. Biophys. Acta2009; 1792:988-97. Wang Y, Guan X, Fok KL, et al. A novel member of the rhomboid family, RHBDD1, regulates BIK-mediated apoptosis. Cell Mol. Life Sci.2008; 65:3822-9.