

IRF2BP2 Antibody
Catalog # ASC11065**Specification**

IRF2BP2 Antibody - Product Information

Application	IHC
Primary Accession	Q7Z5L9
Other Accession	NP_892017 , 116734704
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	IRF2BP2 antibody can be used for detection of IRF2BP2 by immunohistochemistry at 5 µg/mL.

IRF2BP2 Antibody - Additional InformationGene ID **359948****Target/Specificity**

IRF2BP2; At least two isoforms of IRF2BP2 are known to exist; this antibody will detect both.

Reconstitution & Storage

IRF2BP2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

IRF2BP2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

IRF2BP2 Antibody - Protein Information**Name** IRF2BP2**Function**

Acts as a transcriptional corepressor in a IRF2-dependent manner; this repression is not mediated by histone deacetylase activities (PubMed:<[a href="http://www.uniprot.org/citations/12799427"](http://www.uniprot.org/citations/12799427) target="_blank">12799427). Represses the NFAT1-dependent transactivation of NFAT-responsive promoters (PubMed:<[a href="http://www.uniprot.org/citations/21576369"](http://www.uniprot.org/citations/21576369) target="_blank">21576369). Acts as a coactivator of VEGFA expression in cardiac and skeletal muscles (PubMed:<[a href="http://www.uniprot.org/citations/20702774"](http://www.uniprot.org/citations/20702774) target="_blank">20702774). Plays a role in immature B-cell differentiation (PubMed:<[a href="http://www.uniprot.org/citations/27016798"](http://www.uniprot.org/citations/27016798) target="_blank">27016798).

Cellular Location


Cytoplasm. Nucleus.

IRF2BP2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

IRF2BP2 Antibody - Images



Immunohistochemistry of IRF2BP2 in human placenta with IRF2BP2 antibody at 5 µg/mL.

IRF2BP2 Antibody - Background

IRF2BP2 Antibody: IRF2BP2, like the related protein IRF2BP1, is a co-repressor that interacts specifically with the C-terminal repression domain of Interferon Regulatory Factor 2 (IRF2). IRF2BP2 is a direct target gene of p53 and is involved in cell survival during the p53 stress response, able to impede the p53-mediated transactivation of p21 and Bax. IRF2BP2 is also a co-factor of VGLL4 and is required to induce the expression of vascular endothelial growth factor A (VEGF-A) in muscle. It is normally found in the nucleus of skeletal muscle and cardiac cells, but can be found in the cytoplasm during skeletal muscle differentiation.

IRF2BP2 Antibody - References

Childs KS and Goodbourn S. Identification of novel co-repressor molecules for interferon regulatory factor-2. *Nuc. Acids Res.* 2003; 31:3016-26.

Koeppel M, van Heeringen SJ, Smeenk L, et al. The novel p53 target gene IRF2BP2 participates in cell survival during the p53 stress response. *Nuc. Acids Res.* 2009; 37:322-35.

Teng ACT, Kuraitis D, Deeke SA, et al. IRF2BP2 is a skeletal and cardiac muscle-enriched ischemia-inducible activator of VEGFA expression. *FASEB J.* 2010; 24:4825-34.

Teng ACT, Al-montashiri NAM, Cheng BLM, et al. Identification of a phosphorylation-dependent nuclear localization motif in interferon regulatory factor 2 binding protein 2. *PLoS One* 2011; 6:e24100.