

# **LZTR2 Antibody**

Catalog # ASC11074

# **Specification**

# **LZTR2 Antibody - Product Information**

Application
Primary Accession
Other Accession
Reactivity
Host
Clonality
Isotype
Application Notes

WB, IHC-P, IF, E
Q96JE7
NP\_149118, 116235480
Human, Mouse, Rat
Rabbit
Polyclonal
IgG
LZTR2 antibody can be used for detection
of LZTR2 by Western blot at 1 μg/mL.
Antibody can also be used for
immunohistochemistry starting at 5 μg/mL.
For immunofluorescence start at 20 μg/mL.

### **LZTR2 Antibody - Additional Information**

Gene ID Target/Specificity SEC16B; 89866

# **Reconstitution & Storage**

LZTR2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

#### **Precautions**

LZTR2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

#### LZTR2 Antibody - Protein Information

Name SEC16B

Synonyms KIAA1928, LZTR2, RGPR, SEC16S

#### **Function**

Plays a role in the organization of the endoplasmic reticulum exit sites (ERES), also known as transitional endoplasmic reticulum (tER). Required for secretory cargo traffic from the endoplasmic reticulum to the Golgi apparatus (PubMed:<a href="http://www.uniprot.org/citations/17192411" target="\_blank">17192411</a>, PubMed:<a href="http://www.uniprot.org/citations/21768384" target="\_blank">21768384</a>, PubMed:<a href="http://www.uniprot.org/citations/22355596" target="\_blank">22355596</a>). Involved in peroxisome biogenesis. Regulates the transport of peroxisomal biogenesis factors PEX3 and PEX16 from the ER to peroxisomes (PubMed:<a href="http://www.uniprot.org/citations/21768384" target="\_blank">21768384</a>).



### **Cellular Location**

Endoplasmic reticulum membrane; Peripheral membrane protein. Golgi apparatus membrane; Peripheral membrane protein. Note=Localizes to endoplasmic reticulum exit sites (ERES), also known as transitional endoplasmic reticulum (tER).


**Tissue Location** Ubiquitous.

# **LZTR2 Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# **LZTR2 Antibody - Images**




Western blot analysis of LZTR2 in mouse kidney tissue lysate with LZTR2 antibody at 1  $\mu$ g/mL.



Immunohistochemistry of LZTR2 in mouse kidney tissue with LZTR2 antibody at 5 µg/mL.





Immunofluorescence of LZTR2 in mouse kidney tissue with LZTR2 antibody at 20 μg/mL.

# **LZTR2 Antibody - Background**

LZTR2 Antibody: LZTR2, also known as RGPR-p117, is a member of the BTB-kelch superfamily and was initially described as a nuclear factor I (NFI) binding protein and transcriptional regulator of the regucalcin gene. LZTR2 is cytoplasmically localized but is thought to translocate to the nucleus, a process mediated by protein kinase C signaling following hormonal stimulation. Recent evidence has suggested that there is a strong correlation of single nucleotide polymorphisms of LZTR2 with obesity in the Japanese population similar to that seen with the TMEM18 gene and the GNPDA2, BDNF, FAIM2, and MC4R genes with obesity in Caucasian populations, suggesting LZTR2 may play a role in metabolism and obesity risk.

# **LZTR2 Antibody - References**

Misawa H and Yamaguchi M. Molecular cloning and sequencing of the cDNA coding for a novel regucalcin gene promoter region-related protein in rat, mouse and human liver. Int. J. Mol. Med.2001; 8:513-20.

Yamaguchi M. Novel protein RGPR-p117: its role as the regucalcin gene transcription factor. Mol. Cell. Biochem.2009; 327:53-63.

Hotta K, Nakamura M, Nakamura T, et al. Association between obesity and polymorphisms in SEC16B, TMEM18, GNPDA2, BDNF, FAIM2 and MC4R in a Japanese population. J. Hum. Genet.2009; 54:727-31.