

EFHD2 Antibody
Catalog # ASC11090**Specification**

EFHD2 Antibody - Product Information

Application	WB, IHC-P, IF, E
Primary Accession	Q96C19
Other Accession	NP_077305 , 20149675
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	EFHD2 antibody can be used for detection of EFHD2 by Western blot at 1 - 2 µg/mL. Antibody can also be used for immunohistochemistry starting at 5 µg/mL. For immunofluorescence start at 20 µg/mL.

EFHD2 Antibody - Additional Information

Gene ID	79180
Target/Specificity	
EFHD2;	

Reconstitution & Storage

EFHD2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

EFHD2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

EFHD2 Antibody - Protein Information

Name EFHD2

Synonyms SWS1

Function

May regulate B-cell receptor (BCR)-induced immature and primary B-cell apoptosis. Plays a role as negative regulator of the canonical NF-kappa-B-activating branch. Controls spontaneous apoptosis through the regulation of BCL2L1 abundance.

Cellular Location

Membrane raft. Note=In a mouse immature B-cell line WEHI-231.

Tissue Location


Found in lymphocytes; preferentially expressed in CD8+ cells.

EFHD2 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


EFHD2 Antibody - Images



Immunohistochemistry of EFHD2 in human brain tissue with EFHD2 antibody at 5 μ g/mL.



Immunofluorescence of EFHD2 in human brain tissue with EFHD2 antibody at 20 μ g/mL.



Western blot analysis of EFHD2 in mouse brain tissue lysate with EFHD2 antibody at (A) 1 and (B) 2 µg/mL.

EFHD2 Antibody - Background

EFHD2 Antibody: EFHD2, also known as Swiprosin-1 or SWS1, is an EF-hand and coiled-coil-containing adaptor protein that plays a role in lymphocyte physiology. EFHD2 exhibits the highest expression in CD8+ T cells and immature B cells. It provides a membrane scaffold that is required for the Syk-, SLP-65-, and PLCgamma2-dependent B-cell receptor (BCR)-induced calcium flux. EFHD2 may also regulate BCR-induced immature and primary B-cell apoptosis. It controls spontaneous apoptosis through the regulation of BCL2L1 abundance. Also, EFHD2 plays a role as negative regulator of the canonical NF-κB-activating branch.

EFHD2 Antibody - References

Krocze C, Lang C, Brachs S, et al. Swiprosin-1/EFhd2 controls B cell receptor signaling through the assembly of the B cell receptor, Syk, and phospholipase C gamma2 in membrane rafts. *J. Immunol.*2010; 184:3665-76.

Avramidou A, Krocze C, Lang C, et al. The novel adaptor protein Swiprosin-1 enhances BCR signals and contributes to BCR-induced apoptosis. *Cell Death Differ.*2007; 14:1936-47.

Thylur RP, Kiim YD, Kwon MS, et al. Swiprosin-1 is expressed in mast cells and up-regulated through the protein kinase C beta I/eta pathway. *J. Cell Biochem.*2009; 108:705-15.