

**ZC3H12A Antibody**  
**Catalog # ASC11098****Specification****ZC3H12A Antibody - Product Information**

Application	WB, E
Primary Accession	<a href="#">Q5D1E8</a>
Other Accession	<a href="#">NP_079355</a> , <a href="#">156151383</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	Predicted: 66 kDa

Application Notes	Observed: 6 kDa KDa ZC3H12A antibody can be used for detection of ZC3H12A by Western blot at 1 - 2 µg/mL.
-------------------	--

**ZC3H12A Antibody - Additional Information**

Gene ID **80149**

**Target/Specificity**

ZC3H12A; This ZC3H12A antibody is predicted to have no cross-reactivity to other members of the ZC3H12 protein family.

**Reconstitution & Storage**

ZC3H12A antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

**Precautions**

ZC3H12A Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**ZC3H12A Antibody - Protein Information**

**Name** ZC3H12A ([HGNC:26259](#))

**Function**

Endoribonuclease involved in various biological functions such as cellular inflammatory response and immune homeostasis, glial differentiation of neuroprogenitor cells, cell death of cardiomyocytes, adipogenesis and angiogenesis. Functions as an endoribonuclease involved in mRNA decay (PubMed:<a href="http://www.uniprot.org/citations/19909337" target="\_blank">19909337</a>). Modulates the inflammatory response by promoting the degradation of a set of translationally active cytokine-induced inflammation-related mRNAs, such as IL6 and IL12B, during the early phase of inflammation (PubMed:<a href="http://www.uniprot.org/citations/26320658" target="\_blank">26320658</a>). Prevents aberrant T-cell-mediated immune reaction by degradation of multiple mRNAs controlling T-cell

activation, such as those encoding cytokines (IL6 and IL2), cell surface receptors (ICOS, TNFRSF4 and TNFR2) and transcription factor (REL) (By similarity). Inhibits cooperatively with ZC3H12A the differentiation of helper T cells Th17 in lungs. They repress target mRNA encoding the Th17 cell-promoting factors IL6, ICOS, REL, IRF4, NFKBID and NFKBIZ. The cooperation requires RNA-binding by RC3H1 and the nuclease activity of ZC3H12A (By similarity). Together with RC3H1, destabilizes TNFRSF4/OX40 mRNA by binding to the conserved stem loop structure in its 3'UTR (By similarity). Self regulates by destabilizing its own mRNA (By similarity). Cleaves mRNA harboring a stem-loop (SL), often located in their 3'-UTRs, during the early phase of inflammation in a helicase UPF1-dependent manner (PubMed:<a href="http://www.uniprot.org/citations/19909337" target="\_blank">19909337</a>, PubMed:<a href="http://www.uniprot.org/citations/22561375" target="\_blank">22561375</a>, PubMed:<a href="http://www.uniprot.org/citations/26134560" target="\_blank">26134560</a>, PubMed:<a href="http://www.uniprot.org/citations/26320658" target="\_blank">26320658</a>). Plays a role in the inhibition of microRNAs (miRNAs) biogenesis (PubMed:<a href="http://www.uniprot.org/citations/22055188" target="\_blank">22055188</a>). Cleaves the terminal loop of a set of precursor miRNAs (pre-miRNAs) important for the regulation of the inflammatory response leading to their degradation, and thus preventing the biosynthesis of mature miRNAs (PubMed:<a href="http://www.uniprot.org/citations/22055188" target="\_blank">22055188</a>). Also plays a role in promoting angiogenesis in response to inflammatory cytokines by inhibiting the production of antiangiogenic microRNAs via its anti-dicer RNase activity (PubMed:<a href="http://www.uniprot.org/citations/24048733" target="\_blank">24048733</a>). Affects the overall ubiquitination of cellular proteins (By similarity). Positively regulates deubiquitinase activity promoting the cleavage at 'Lys-48'- and 'Lys-63'-linked polyubiquitin chains on TNF receptor-associated factors (TRAFs), preventing JNK and NF-kappa-B signaling pathway activation, and hence negatively regulating macrophage-mediated inflammatory response and immune homeostasis (By similarity). Also induces deubiquitination of the transcription factor HIF1A, probably leading to its stabilization and nuclear import, thereby positively regulating the expression of proangiogenic HIF1A-targeted genes (PubMed:<a href="http://www.uniprot.org/citations/24048733" target="\_blank">24048733</a>). Involved in a TANK-dependent negative feedback response to attenuate NF-kappaB activation through the deubiquitination of IKBKG or TRAF6 in response to interleukin-1-beta (IL1B) stimulation or upon DNA damage (PubMed:<a href="http://www.uniprot.org/citations/25861989" target="\_blank">25861989</a>). Prevents stress granule (SGs) formation and promotes macrophage apoptosis under stress conditions, including arsenite- induced oxidative stress, heat shock and energy deprivation (By similarity). Plays a role in the regulation of macrophage polarization; promotes IL4-induced polarization of macrophages M1 into anti- inflammatory M2 state (By similarity). May also act as a transcription factor that regulates the expression of multiple genes involved in inflammatory response, angiogenesis, adipogenesis and apoptosis (PubMed:<a href="http://www.uniprot.org/citations/16574901" target="\_blank">16574901</a>, PubMed:<a href="http://www.uniprot.org/citations/18364357" target="\_blank">18364357</a>). Functions as a positive regulator of glial differentiation of neuroprogenitor cells through an amyloid precursor protein (APP)-dependent signaling pathway (PubMed:<a href="http://www.uniprot.org/citations/19185603" target="\_blank">19185603</a>). Attenuates septic myocardial contractile dysfunction in response to lipopolysaccharide (LPS) by reducing I-kappa-B-kinase (IKK)-mediated NF-kappa-B activation, and hence myocardial pro-inflammatory cytokine production (By similarity).

### Cellular Location

Nucleus. Cytoplasm. Cytoplasm, P-body. Rough endoplasmic reticulum membrane {ECO:0000250|UniProtKB:Q5D1E7}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q5D1E7}; Cytoplasmic side {ECO:0000250|UniProtKB:Q5D1E7}. Cytoplasmic granule {ECO:0000250|UniProtKB:Q5D1E7}. Note=Predominantly localized in the cytoplasm. Colocalizes with GW182 on many granule-like structures, probably corresponding to cytoplasmic GW bodies (GWBs), also called processing bodies (P bodies). Colocalizes with calnexin on the surface of the rough endoplasmic reticulum (RER) membrane and with translationally active polysomes (By similarity). Colocalizes with ZC3H12D in cytoplasmic mRNA processing P-body, also known as GW bodies (GWBs) (PubMed:22055188, PubMed:26134560)

### Tissue Location


Expressed in heart, placenta, spleen, kidney, liver and lung (PubMed:19909337). Expressed in leukocytes (PubMed:19909337) Expressed in monocyte (PubMed:16574901).

### ZC3H12A Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### ZC3H12A Antibody - Images



Western blot analysis of ZC3H12A in K562 cell lysate with ZC3H12A antibody at 1  $\mu$ g/mL in (A) the absence and (B) the presence of blocking peptide.

### ZC3H12A Antibody - Background

**ZC3H12A Antibody:** ZC3H12A, also known as MCPIP, is an essential member of a family of novel CCH-zinc finger proteins that regulate macrophage activation and may be involved in host immunity and inflammatory diseases. ZC3H12A has RNase activity that prevents some immune disorders by directly affecting the mRNA stability of interleukins such as IL-6 and IL12p40. Mice lacking the ZC3H12A gene suffered from severe anemia, and most dies within 12 weeks. Overexpression of ZC3H12A causes ER stress and induces a number of genes involved in apoptosis and autophagy, including JNK, PUMA, and beclin-1 in cardiac myoblasts, leading to cell death. ZC3H12A can also induce adipogenesis in 3T3-L1 pre-adipocytes in the absence of PPAPgamma, demonstrating the complex roles ZC3H12A plays.

### ZC3H12A Antibody - References

Zhou L, Azfer A, Niu J, et al. Monocyte chemoattractant protein-1 induces a novel transcription

factor that causes cardiac myocyte apoptosis and ventricular dysfunction. Circ. Res. 2006; 98:1177-85.

Liang J, Wang J, Azfer A, et al. A novel CCCH-zinc finger protein family regulates proinflammatory activation of macrophages. J. Biol. Chem. 2008; 283:6337-46.

Matsushita K, Takeuchi O, Standley DM, et al. Zc3h12a is an RNase essential for controlling immune responses by regulating mRNA decay. Nature 2009; 458:1185-90.

Younce CW and Kolattukudy PE. MCP-1 causes cardiomyoblast death via autophagy resulting from ER stress caused by oxidative stress generated by inducing a novel zinc-finger protein, MCPIP. Biochem. J. 2010; 426:43-53.