

SCRN2 Antibody

Catalog # ASC11169

Specification

SCRN2 Antibody - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Isotype **Application Notes**

WB, ICC, IF **Q96FV2**

EAW94970, 119615376 Human, Mouse, Rat

Rabbit Polyclonal

laG

SCRN2 antibody can be used for detection of SCRN2 by Western blot at 0.5 - 1 μg/mL.

Antibody can also be used for

immunocytochemistry starting at 20

μg/mL. For immunofluorescence start at 20

μg/mL.

SCRN2 Antibody - Additional Information

90507 Gene ID

Target/Specificity

SCRN2:

Reconstitution & Storage

SCRN2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

SCRN2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

SCRN2 Antibody - Protein Information

Name SCRN2

SCRN2 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence




- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture


SCRN2 Antibody - Images



Western blot analysis of SCRN2 in 293 cell tissue lysate with SCRN2 antibody at (A) 0.5 and (B) 1 $\mu g/mL$.



Immunocytochemistry of SCRN2 in 293 cells with SCRN2 antibody at 20 μg/mL.



Immunofluorescence of SCRN2 in 293 cells with SCRN2 antibody at 20 $\mu g/mL$.

SCRN2 Antibody - Background

SCRN2 Antibody: SCRN2 is a member of the secernin family of proteins which includes the





Tel: 858.875.1900 Fax: 858.875.1999

homologous SCRN1, a protein that was first identified as being involved in the regulation of exocytosis from peritoneal mast cells. Little is known of SCRN2, but studies have shown that SCRN1 may possess epitopes that function as tumor-associated antigens in gastric cancers and increased SCRN1 expression in patients with colorectal cancer correlated with poor prognosis, suggesting that SCRN2 may also be involved in protein secretion or the regulation of cell growth.

SCRN2 Antibody - References

Way G, Morrice N, Smythe C, et al. Purification and identification of Secernin, a novel cytosolic protein that regulates exocytosis in mast cells. Mol. Biol. Cell2002; 13:3344-54. Suda T, Tsunoda T, Uchida N, et al. Identification of secernin 1 as a novel immunotherapy target for gastric cancer using the expression profiles of cDNA microarray. Cancer Sci.2006; 97:411-9. Miyoshi N, Ishii H, Mimori K, et al. SCRN1 is a novel marker for prognosis in colorectal cancer. J. Surg. Oncol.2010; 101:156-9.