

# NOD3 Antibody

Catalog # ASC11189

### Specification

### NOD3 Antibody - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Isotype Application Notes

WB, IF, E <u>Q7RTR2</u> <u>EAW85351</u>, <u>119605757</u> Human, Mouse, Rat Rabbit Polyclonal IgG NOD3 antibody can be used for detection of NOD3 by Western blot at 1 - 2 μg/mL. Antibody can also be used for immunofluorescence starting at 20 μg/mL. For immunofluorescence start at 20 μg/mL.

## NOD3 Antibody - Additional Information

Gene ID Target/Specificity NLRC3;

197358

#### **Reconstitution & Storage**

NOD3 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

**Precautions** NOD3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

### **NOD3 Antibody - Protein Information**

Name NLRC3

Synonyms NOD3

#### Function

Negative regulator of the innate immune response (PubMed:<a

href="http://www.uniprot.org/citations/15705585" target="\_blank">15705585</a>, PubMed:<a href="http://www.uniprot.org/citations/22863753" target="\_blank">22863753</a>, PubMed:<a href="http://www.uniprot.org/citations/25277106" target="\_blank">22863753</a>, PubMed:<a href="http://www.uniprot.org/citations/25277106" target="\_blank">25277106</a>). Attenuates signaling pathways activated by Toll-like receptors (TLRs) and the DNA sensor STING/TMEM173 in response to pathogen-associated molecular patterns, such as intracellular poly(dA:dT), but not poly(I:C), or in response to DNA virus infection, including that of Herpes simplex virus 1 (HSV1) (By similarity) (PubMed:<a href="http://www.uniprot.org/citations/22863753" target=" blank">22863753</a>). May affect TLR4 signaling by acting at the level of TRAF6



ubiquitination, decreasing the activating 'Lys-63'-linked ubiquitination and leaving unchanged the degradative 'Lys-48'-linked ubiquitination (PubMed:<a

href="http://www.uniprot.org/citations/22863753" target="\_blank">22863753</a>). Inhibits the PI3K-AKT-mTOR pathway possibly by directly interacting with the posphatidylinositol 3-kinase regulatory subunit p85 (PIK3R1/PIK3R2) and disrupting the association between PIK3R1/PIK3R2 and the catalytic subunit p110 (PIK3CA/PIK3CB/PIK3CD) and reducing PIK3R1/PIK3R2 activation. Via its regulation of the PI3K-AKT-mTOR pathway, controls cell proliferation, predominantly in intestinal epithelial cells (By similarity). May also affect NOD1- or NOD2-mediated NF-kappa-B activation (PubMed:<a href="http://www.uniprot.org/citations/25277106" target="\_blank">25277106</a>). Might also affect the inflammatory response by preventing NLRP3 inflammasome formation, CASP1 cleavage and IL1B maturation (PubMed:<a href="http://www.uniprot.org/citations/25277106"/a>).

Cellular Location Cytoplasm

### **NOD3 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

#### NOD3 Antibody - Images



Western blot analysis of NOD3 in 3T3 cell lysate with NOD3 antibody at (A) 1 and (B) 2 µg/mL.





Immunofluorescence of NOD3 in Jurkat cells with NOD3 antibody at 20 µg/mL.

## NOD3 Antibody - Background

NOD3 Antibody: NOD3 is a member of the NOD (nucleotide-binding oligomerization domain) family, a group of proteins that are involved in innate immune defense. NOD3 also known as NLR family CARD containing 3 (NLRC3), is predominantly expressed in the immune system, particularly in T lymphocytes, and its expression is strongly down-regulated following stimulation of the T-cell receptor complex and CD28, suggesting that NOD3 plays a role in attenuating the activation of T cells. NOD3 inhibits NF-kappaB, AP-1 and NFAT transcriptional activation in Jurkat T cells downstream of CD3/CD28 stimulation or treatment with PMA/ionomycin and decreases IL-2 and CD25 mRNA induction in activated cells.

### **NOD3 Antibody - References**

Kufer TA, Banks DJ, and Philpott DJ. Innate immune sensing of microbes by Nod proteins. Ann. NY Acad. Sci.2006; 1072:19-27.

Conti BJ, Davis BK, Zhang J, et al. CATERPILLER 16.2 (CLR16.2), a novel NBD/LRR family member that negatively regulates T cell function. J. Biol. Chem.2005; 280:18375-85.