

# ZNF521 Antibody

Catalog # ASC11241

#### Specification

## ZNF521 Antibody - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Isotype Application Notes IHC-P, E <u>O96K83</u> NP\_056276, 24308069 Human Rabbit Polyclonal IgG ZNF521 antibody can be used for detection of ZNF521 by immunohistochemistry at 5 μg/mL.

### ZNF521 Antibody - Additional Information

Gene ID Target/Specificity ZNF521; ZNF521 antibody is human specific.

**Reconstitution & Storage** 

ZNF521 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

25925

#### Precautions

ZNF521 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

#### **ZNF521 Antibody - Protein Information**

Name ZNF521

Synonyms EHZF, LIP3

Function

Transcription factor that can both act as an activator or a repressor depending on the context. Involved in BMP signaling and in the regulation of the immature compartment of the hematopoietic system. Associates with SMADs in response to BMP2 leading to activate transcription of BMP target genes. Acts as a transcriptional repressor via its interaction with EBF1, a transcription factor involved specification of B-cell lineage; this interaction preventing EBF1 to bind DNA and activate target genes.

**Cellular Location** Nucleus.

**Tissue Location** 



Predominantly expressed in hematopoietic cells. Present in organs and tissues that contain stem and progenitor cells, myeloid and/or lymphoid: placenta, spleen, lymph nodes, thymus, bone marrow and fetal liver. Within the hematopoietic system, it is abundant in CD34(+) cells but undetectable in mature peripheral blood leukocytes, and its levels rapidly decrease during the differentiation of CD34(+) cells in response to hemopoietins

#### **ZNF521 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

## ZNF521 Antibody - Images



Immunohistochemistry of ZNF521 in human kidney tissue with ZNF521 antibody at 5 µg/mL.

#### ZNF521 Antibody - Background

ZNF521 Antibody: The zinc finger protein 521 (ZNF521) is a transcription factor containing an N-terminal transcriptional repressor motif and 30 zinc finger domains. It plays a role in both erythroid cell and osteoblast differentiation during development, inhibiting the activities of early B-cell factor 1 (EBF1) in erythroid cells and Runx2 in osteoblast precursors. ZFP521 binds to both Runx2 and histone deacetylase 3 (HDAC3), promotes their association and antagonizes Runx2 transcriptional activity in a HDAC3-dependent manner, thereby regulating osteoblast differentiation, skeletal development, and bone homeostasis.

#### ZNF521 Antibody - References

Warming S, Liu P, Suzuki T, et al. Evi3, a common retroviral integration site in murine B-cell lymphoma, encodes a EBFAZ-related Kruppel-like zinc finger protein. Blood 2003; 101:1934-40. Matsubara E, Sakai I, Yamanouchi J, et al. The role of zinc finger protein 521/early hematopoietic zinc finger protein in erythroid cell differentiation. J. Biol. Chem. 2009; 284:3480-7. Wu M, Hesse E, Morvan F, et al. Zfp521 antagonizes Runx2, delays osteoblast differentiation in vitro, and promotes bone formation in vivo. Bone 2009; 44:528-36. Hesse E, Saito H, Kiviranta R, et al. Zfp521 controls bone mass by HDAC3-dependent attenuation of



Runx2 activity. J. Cell Biol. 2010; 191:1271-83.