

LRRTM1 Antibody
Catalog # ASC11274**Specification**

LRRTM1 Antibody - Product Information

Application	WB, IHC, IF
Primary Accession	Q86UE6
Other Accession	AAH45113 , 86990456
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	LRRTM1 antibody can be used for detection of LRRTM1 by Western blot at 1 µg/mL. Antibody can also be used for immunohistochemistry starting at 5 µg/mL. For immunofluorescence start at 20 µg/mL.

LRRTM1 Antibody - Additional InformationGene ID **347730****Target/Specificity**

LRRTM1; LRRTM1 antibody is predicted to not cross-react with other LRRTM family members.

Reconstitution & Storage

LRRTM1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

LRRTM1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

LRRTM1 Antibody - Protein Information**Name** LRRTM1**Function**

Exhibits strong synaptogenic activity, restricted to excitatory presynaptic differentiation, acting at both pre- and postsynaptic level.

Cellular Location

Cell membrane; Single-pass type I membrane protein. Postsynaptic cell membrane; Single-pass type I membrane protein

Tissue Location


Predominantly expressed in forebrain regions including thalamus and cerebral cortex.

LRRTM1 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


LRRTM1 Antibody - Images



Western blot analysis of LRRTM1 in mouse brain tissue lysate with LRRTM1 antibody at 1 μ g/mL in (A) the absence and (B) the presence of blocking peptide.



Immunohistochemistry of LRRTM1 in human brain tissue with LRRTM1 antibody at 5 μ g/mL.



Immunofluorescence of LRRTM1 in human brain tissue with LRRTM1 antibody at 20 µg/mL.

LRRTM1 Antibody - Background

LRRTM1 Antibody: The Leucine-rich repeat transmembrane neuronal proteins (LRRTMs) are differentially expressed in the nervous system and were recently found to instruct presynaptic and mediate postsynaptic glutamatergic differentiation, with LRRTM1 and LRRTM2 most potent at inducing presynaptic differentiation. Each LRRTM protein is a type I transmembrane containing ten extracellular leucine-rich repeats and a short intracellular tail and has a developmentally regulated pattern distinct from all others. LRRTM1 is a maternally suppressed gene that is associated paternally with handedness and schizophrenia.

LRRTM1 Antibody - References

Lauren J, Airaksinen MS, Saarma M, et al. A novel gene family encoding leucine-rich repeat transmembrane protein differentially expressed in the nervous system. *Genomics* 2003; 81:411-21.
Linhoff MW, Lauren J, Cassidy RM, et al. An unbiased expression screen for synaptogenic proteins identifies the LRRTM protein family as synaptic organizers. *Neuron* 2009; 61:734-49.
Siddiqui TJ, Pancaroglu R, Kang Y, et al. LRRTMs and neuroligins bind neurexins with a differential code to cooperate in glutamate synapse development. *J. Neurosci.* 2010; 30:7495-506.
Haines BP and Rigby PW. Developmentally regulated expression of the LRRTM gene family during mid-gestation mouse embryogenesis. *Gene Expr. Patterns* 2007; 7:23-9.