

EIG121 Antibody
Catalog # ASC11360**Specification**

EIG121 Antibody - Product Information

Application	WB, IF, ICC, E
Primary Accession	Q6UXG2
Other Accession	NP_065826 , 38569482
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application Notes	EIG121 antibody can be used for detection of EIG121 by Western blot at 1 µg/mL. Antibody can also be used for immunocytochemistry starting at 5 µg/mL. For immunofluorescence start at 20 µg/mL.

EIG121 Antibody - Additional InformationGene ID **1957****Target/Specificity**

EIG1; At least four isoforms of EIG121 known to exist. EIG121 antibody is predicted to not cross-react with other UPF0577 protein family members.

Reconstitution & Storage

EIG121 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

EIG121 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

EIG121 Antibody - Protein InformationName ELAPOR1 ([HGNC:29618](#))**Function**

May protect cells from cell death by inducing cytosolic vacuolization and up-regulating the autophagy pathway (PubMed:<[a href="http://www.uniprot.org/citations/21072319"](http://www.uniprot.org/citations/21072319)>21072319). May play a role in apoptosis and cell proliferation through its interaction with HSPA5 (PubMed:<[a href="http://www.uniprot.org/citations/26045166"](http://www.uniprot.org/citations/26045166)>26045166).

Cellular Location

Cell membrane; Single-pass type I membrane protein. Late endosome membrane; Single-pass type I membrane protein. Golgi apparatus, trans-Golgi network membrane; Single-pass type I membrane protein. Lysosome membrane; Single-pass membrane protein. Endoplasmic reticulum

membrane; Single-pass type I membrane protein

Tissue Location


Expressed in normal endometrium but overexpressed in endometroid tumors.

EIG121 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


EIG121 Antibody - Images



Western blot analysis of EIG121 in MCF7 cell lysate with EIG121 antibody at 1 μ g/mL in (A) the absence and (B) the presence of blocking peptide.



Immunocytochemistry of EIG121 in MCF7 cells with EIG121 antibody at 5 μ g/mL.



Immunofluorescence of EIG121 in MCF7 cells with EIG121 antibody at 20 µg/mL.

EIG121 Antibody - Background

EIG121 Antibody: EIG121 (Estrogen-induced gene 121 protein) is thought to play a role as a marker of hyperestrogenic state and estrogen-related type I endometrial carcinoma. It belongs to the UPF0577 family. It is a 1013 amino acid single-pass transmembrane protein that, though expressed in normal endometrium, is overexpressed in endometrial tumors. As an evolutionarily conserved gene, EIG121 is also expressed during early xenopus development, showing maximum expression at the gastrula stage.

EIG121 Antibody - References

Deng L, Broaddus RR, McCampbell A, et al. Identification of a novel estrogen-regulated gene, EIG121, induced by hormone replacement therapy and differentially expressed in type I and type II endometrial cancer. Clin. Cancer Res. 2005; 11:8258-64.

Deng L, Feng J, Broaddus RR. The novel estrogen-induced gene EIG121 regulates autophagy and promotes cell survival under stress. Cell Death Dis. 2010; 1:e32.

Araki T, Kusakabe M and Nishida E. Expression of estrogen induced gene 121-like (EIG121L) during early Xenopus development. Gene Expr. Patterns 2007; 7:666-71.