

CBX4 Antibody
Catalog # ASC11535**Specification**

CBX4 Antibody - Product Information

Application	WB, IF, E
Primary Accession	O00257
Other Accession	NP_003646 , 55770830
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	62 kDa KDa
Application Notes	CBX4 antibody can be used for detection of CBX4 by Western blot at 1 µg/mL. For immunofluorescence start at 20 µg/mL.

CBX4 Antibody - Additional InformationGene ID **8535****Target/Specificity**

CBX4; Two alternatively spliced transcript variants have been observed.

Reconstitution & Storage

CBX4 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

CBX4 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

CBX4 Antibody - Protein Information**Name** CBX4**Function**

E3 SUMO-protein ligase that catalyzes sumoylation of target proteins by promoting the transfer of SUMO from the E2 enzyme to the substrate (PubMed:[12679040](http://www.uniprot.org/citations/12679040), PubMed:[22825850](http://www.uniprot.org/citations/22825850)). Involved in the sumoylation of HNRNPK, a p53/TP53 transcriptional coactivator, hence indirectly regulates p53/TP53 transcriptional activation resulting in p21/CDKN1A expression. Monosumoylates ZNF131 (PubMed:[22825850](http://www.uniprot.org/citations/22825850)).

Cellular Location

Nucleus. Nucleus speckle. Note=Localization to nuclear polycomb bodies is required for ZNF131 sumoylation (PubMed:22467880). Localized in distinct foci on chromatin (PubMed:18927235)


Tissue Location

Ubiquitous.


CBX4 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CBX4 Antibody - Images

Western blot analysis of CBX4 in human brain tissue lysate with CBX4 antibody at 1 µg/ml in (A) the absence and (B) the presence of blocking peptide.



Immunofluorescence of CBX4 in human brain tissue with CBX4 antibody at 20 µg/mL.

CBX4 Antibody - Background

CBX4 Antibody: Polycomb group (PcG) proteins are chromatin-associated proteins which are important for embryonic and adult stem cell self-renewal and maintenance. At least two distinct human PcG complexes have been identified. Polycomb repressive complex I (PRC1) contains CBX4/Pc2, HPH1 and RING domain-containing proteins (RING1, RING2 and BMI1). CBX4 (Chromobox protein homolog 4 or PC2), which functions as a transcriptional suppressor, is a candidate of KyoT2-binding proteins. It is a E3 SUMO-protein ligase which facilitates SUMO1 conjugation. Depletion of CBX4 results in decreased cellular resistance to ionizing radiation. CBX4 is directly involved in the cellular response to DNA damage.

CBX4 Antibody - References

Simon JA and Tamkun JW. Programming off and on states in chromatin: mechanisms of Polycomb and trithorax group complexes. *Curr. Opin. Genet. Dev.* 2002;12:210-8.

Gieni RS and Hendzel MJ. Polycomb group protein gene silencing, non-coding RNA, stem cells, and cancer. *Biochem. Cell Biol.* 2009; 87:711-46.

Luis NM, Morey L, Mejetta S, et al. Regulation of human epidermal stem cell proliferation and senescence requires polycomb-dependent and -independent functions of Cbx4. *Cell. Stem Cell* 2011; 9:233-46.

Ismail IH, Gagné JP, Caron MC, et al. CBX4-mediated SUMO modification regulates BMI1 recruitment at sites of DNA damage. *Nucleic Acids Res.* 2012 Mar 8.