

## **JAKMIP1 Antibody**

Catalog # ASC11593

### **Specification**

# JAKMIP1 Antibody - Product Information

Application
Primary Accession
Other Accession
Reactivity
Host

Clonality Isotype

Calculated MW

**Application Notes** 

WB, E 096N16

NP\_653321, 21389505 Human, Mouse, Rat

Rabbit Polyclonal

IgG

Predicted: 69 kDa KDa

JAKMIP1 antibody can be used for

detection of JAKMIP1 by Western blot at 1 -

2 μg/mL.

## **JAKMIP1** Antibody - Additional Information

Gene ID **152789** 

**Target/Specificity** 

JAKMIP1; Multiple isoforms of JAKMIP1 are known to exist. JAKMIP1 antibody is predicted to not cross-react with JAKMIP2.

### **Reconstitution & Storage**

JAKMIP1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.

## **Precautions**

JAKMIP1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

# JAKMIP1 Antibody - Protein Information

### Name JAKMIP1

Synonyms GABABRBP, JAMIP1, MARLIN1

#### **Eunction**

Associates with microtubules and may play a role in the microtubule-dependent transport of the GABA-B receptor. May play a role in JAK1 signaling and regulate microtubule cytoskeleton rearrangements.

## **Cellular Location**

Cytoplasm, cytoskeleton. Membrane; Peripheral membrane protein. Note=Colocalizes with the microtubule network Localizes to the cell body and neurites of hippocampal neurons where it accumulates in granules. Localizes to the tail and to a lower extent to the head of sperm cells

#### **Tissue Location**

Predominantly expressed in neural tissues and lymphoid cells (at protein level). Isoform 2, isoform



Tel: 858.875.1900 Fax: 858.875.1999


3 and isoform 4 are specifically expressed in brain and retina. Isoform 1 and isoform 5 are also detected in liver, lung and skeletal muscle. Also detected in testis and to a lower extent spleen and intestine

### **JAKMIP1 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

## **JAKMIP1 Antibody - Images**



Western blot analysis of JAKMIP1 in rat brain tissue lysate with JAKMIP1 antibody at (A) 1 and (B)  $2 \mu g/mL$ 

### JAKMIP1 Antibody - Background


JAKMIP1 Antibody: JAKMIP1, also known as Marlin-1, is a highly-conserved vertebrate-specific microtubule-associated protein that plays a role in JAK1 signaling and regulate microtubule cytoskeleton rearrangements. JAKMIP1 is expressed predominantly in neuronal and lymphoid cells and colocalizes with microtubules. Multiple isoforms of JAKMIP1 are known to exist: isoforms 1 and 5 are also expressed in liver, lung and skeletal muscle, while isoforms 2-4 are specifically expressed in brain and retina. JAKMIP1 participates in polarized secretion in lymphocytes and it has been suggested to be a novel effector memory gene restraining T cell-mediated cytotoxicity. Dysregulation of JAKMIP1 gene expression is associated with genetic conditions linked to autism in humans.

## **JAKMIP1 Antibody - References**

Costa V, Conte I, Ziviello C, et al. Identification and expression analysis of novel Jakmip1 transcripts. Gene 2007; 402:1-8.

Vidal RL, Fuentes P, Valenzuela JI, et al. RNA interference of Marlin-1/Jakmip1 results in abnormal







morphogenesis and migration of cortical pyramidal neurons. Mol. Cell Neurosci. 2012; epub. Libri V, Schulte D, Van Stijn A et al. Jakmip1 is expressed upon T cell differentiation and has an inhibitory function in cytotoxic T lymphocytes. J. Immunol. 2008; 181:5847-56 Bill BR, Geschwind DH. Genetic advances in autism: heterogeneity and convergence on shared pathways. Curr. Opin. Genet. Dev. 2009; 19:271-8.