

Alpha-tubulin Antibody
Catalog # ASC11595**Specification**

Alpha-tubulin Antibody - Product Information

Application	WB, E
Primary Accession	Q13748
Other Accession	NP_005992 , 17921993
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	Predicted: 50 kDa KDa
Application Notes	Tubulin antibody can be used for detection of Tubulin by Western blot at 0.5 and 1 µg/mL.

Alpha-tubulin Antibody - Additional Information

Gene ID	7278
Target/Specificity	
TUBA3C;	

Reconstitution & Storage

Alpha-tubulin antibody can be stored at 4°C for three months and -20°C, stable for up to one year.

Precautions


Alpha-tubulin Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Alpha-tubulin Antibody - Protein Information**Alpha-tubulin Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Alpha-tubulin Antibody - Images



Western blot analysis of Tubulin in mouse brain tissue lysate with Tubulin antibody at (A) 0.5 and (B) 1 μ g/mL.

Alpha-tubulin Antibody - Background

Alpha-tubulin Antibody: Alpha-tubulin belongs to the tubulin superfamily, which is composed of six distinct families. Along with beta-tubulins, alpha-tubulins are the major components of microtubules. These microtubules are involved in a wide variety of cellular activities ranging from mitosis and transport events to cell movement and the maintenance of cell shape. Alpha- and beta-tubulin dimers are assembled to 13 protofilaments that form a microtubule of 22-nm diameter. Tyrosine ligase adds a C-terminal tyrosine to monomeric alpha-tubulin. Assembled microtubules can again be detyrosinated by a cytoskeleton-associated carboxypeptidase. Another post-translational modification of detyrosinated alpha-tubulin is C-terminal polyglutamylation, which is characteristic of microtubules in neuronal cells and the mitotic spindle. Like GAPDH and beta-Actin, this antibody makes an excellent loading control in immunoblots.

Alpha-tubulin Antibody - References

McKean PG, Vaughan S, and Gull K. The extended tubulin family. *J. Cell Sci.* 2001; 114:2723-33.
Barra HA, Arce CA, and Argarana CE. Posttranslational tyrosination/detyrosination of tubulin. *Mol. Neurobiol.* 1988; 2:133-53.
Fukushima N, Furuta D, Hidaka Y, et al. Post-translational modifications of tubulin in the nervous system. *J. Neurochem.* 2009; 109:683-693.