

PIBF1 Antibody
Catalog # ASC11618**Specification**

PIBF1 Antibody - Product Information

Application	WB, IHC-P, IF, E
Primary Accession	Q8WXW3
Other Accession	NP_006337 , 5576958
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	83 kDa KDa
Application Notes	PIBF1 antibody can be used for detection of PIBF1 by Western blot at 1 - 2 µg/mL.

PIBF1 Antibody - Additional InformationGene ID **10464****Target/Specificity**

PIBF1; Multiple isoforms of PIBF1 exists as a result of alternative splicing event.

Reconstitution & Storage

PIBF1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.

Precautions

PIBF1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

PIBF1 Antibody - Protein Information**Name** PIBF1**Synonyms** C13orf24, PIBF**Function**

Plays a role in ciliogenesis.

Cellular Location

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm. Secreted
Note=In progesterone-treated astrocytoma cells a 57 kDa protein and isoform 1 (90 kDa) have been described, both being located in the intracellular medium and secreted. Respective predominant forms are isoform 1 in the intracellular and the 57 kDa protein in the extracellular medium (PubMed:25218441). [Isoform 4]: Secreted Note=Secreted by progesterone-treated lymphocytes (PubMed:14634107)

Tissue Location

Expressed at highest levels in testis. Moderate expression is detected in spleen, thymus, prostate, ovary, small intestine, and colon (PubMed:11935316). Expressed in the first trimester pregnancy


decidua (PubMed:12516630). Localized to extravillous cytotrophoblast (at protein level). Also found in syncytiotrophoblast and part of the villous cytotrophoblast. Isoform 1 is expressed in first trimester and term villous trophoblast; trophoblast cells can additionally express other isoforms (PubMed:18817979). Overexpressed in solid tumors from stomach and uterus and in cells from ovary, cervical, breast, lymphoma and leukemia cancer (PubMed:25218441).

PIBF1 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


PIBF1 Antibody - Images



Western blot analysis of PIBF1 in human placenta tissue lysate with PIBF1 antibody at (A) 1 and (B) 2 µg/mL



Immunohistochemistry of PIBF in spleen tissue with PIBF antibody at 5 µg/ml.



Immunofluorescence of PIBF in human spleen tissue with PIBF antibody at 20 µg/ml.

PIBF1 Antibody - Background

PIBF1 Antibody: PIBF1 is synthesized during pregnancy in response to progesterone by T lymphocytes. PIBF1 inhibits arachidonic acid release, controls NK activity, and modifies the cytokine balance exerting an anti-abortion effect. It contains a leucine zipper motif, a basic zipper sequence, a PEST sequence, a nuclear localization signal, an ER membrane retention signal and N-glycosylation and phosphorylation sites. PIBF1 is significantly higher in healthy pregnant women than in women at risk for premature pregnancy termination. Full-length PIBF1 is associated with the nucleus and functions as a transcription factor, whereas secretion of shorter forms which may act as cytokines is induced by activation of the cell.

PIBF1 Antibody - References

Laskarin G, Tokmadzic VS, Strbo N, et al. Progesterone induced blocking factor (PIBF) mediates progesterone induced suppression of decidual lymphocyte cytotoxicity. *Am. J. Reprod. Immunol.* 2002; 48:201-9.

Lachmann M, Gelbmann D, Kalman E, et al. PIBF (progesterone induced blocking factor) is overexpressed in highly proliferating cells and associated with the centrosome. *Int. J. Cancer.* 2004; 112:51-60

Polgar B, Kispal G, Lachmann M, et al. Molecular cloning and immunologic characterization of a novel cDNA coding for progesterone-induced blocking factor. *J. Immunol.* 2003; 171:5956-63.

Raghupathy R, Al-Mutawa E, Al-Azemi M, et al. Progesterone-induced blocking factor (PIBF) modulates cytokine production by lymphocytes from women with recurrent miscarriage or preterm delivery. *J. Reprod. Immunol.* 2009; 80:91-9.