

GOLGA5 Antibody

Catalog # ASC11668

Specification

GOLGA5 Antibody - Product Information

Application WB, IHC, IF Primary Accession O8TBA6

Other Accession
Reactivity
Host
Rebit

Clonality Polyclonal Isotype IgG

Calculated MW Predicted: 64, 80 kDa

Observed: 65, 78 kDa KDa

Application Notes GOLGA5 antibody can be used for

detection of GOLGA5 by Western blot at 1 - 2 μ g/mL. Antibody can also be used for Immunohistochemistry starting at 5 μ g/mL. For immunofluorescence start at 20 μ g/mL.

GOLGA5 Antibody - Additional Information

Gene ID 9950

Target/Specificity

GOLGA5; GOLGA5 antibody is human, mouse and rat reactive. Two isoforms of GOLGA5 exist due to alternative splicing events.

Reconstitution & Storage

GOLGA5 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.

Precautions

GOLGA5 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

GOLGA5 Antibody - Protein Information

Name GOLGA5

Synonyms RETII, RFG5

Function

Involved in maintaining Golgi structure. Stimulates the formation of Golgi stacks and ribbons. Involved in intra-Golgi retrograde transport.

Cellular Location

Golgi apparatus membrane; Single-pass type IV membrane protein. Note=Found throughout the Golgi, both on cisternae and, at higher abundance, on the tubulo-vesicular structures of the cis-Golgi network



Tissue Location


Ubiquitous. Highly expressed in seminiferous tubules and Leydig cells in testis, and detected at much lower levels in the other tissues tested. Expression is very low or not detectable in spermatozoa.

GOLGA5 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

GOLGA5 Antibody - Images




Western blot analysis of GOLGA5 in human testis tissue lysate with GOLGA5 antibody at 1 µg/mL.



Immunohistochemistry of GOLGA5 in human testis tissue with GOLGA5 antibody at 5 μg/mL.





Immunofluorescence of GOLGA5 in human testis tissue with GOLGA5 antibody at 20 µg/mL.

GOLGA5 Antibody - Background

GOLGA5 Antibody: GOLGA5 (Golgi autoantigen, subfamily A, member 5), also known as GOLGA84, RFG5 (RET-fused gene 5 protein) or GOLIM5, is a ubiquitously expressed single-pass type II coiled-coil membrane protein that is involved in maintaining Golgi structure (1). It is localized to the Golgi apparatus and predominantly found on membranes at the cis side of the Golgi stack (1,2). As a mitotic phosphoprotein, GOLGA5 binds to active Rab 1 and associates with CASP (an isoform of CDP) in a golgin-tethering complex that is believed to play a role in intra-Golgi retrograde transport (3,4). Defects in GOLGA5 are a cause of thyroid papillary carcinoma (TPC) (5).

GOLGA5 Antibody - References

Bascom RA, Srinivasan S, Nussbaum RL. Identification and characterization of golgin 84, a novel Golgi integral membrane protein with a cytoplasmic coiled-coil domain. J. Biol. Chem. 1999; 274:2953-62.

Diao A, Rahman D, Pappin DJ, et al. 2003. The coiled-coil membrane protein golgin 84 is a novel Rab effector required for Golgi ribbon formation. J. Cell Biol. 2003; 160:201-12.

Malsam J, Satoh A, Pelletier L, et al. Golgin tethers define subpopulations of COPI vesicles. Science 2005; 307:1095-8.

Sohda M, Misumi Y, Yamamoto A, et al. Interaction of Golgin-84 with the COG complex mediates the intra-Golgi retrograde transport. Traffic 2010;11:1552-66.