

**ERAP2 Antibody**  
**Catalog # ASC11695****Specification**

---

**ERAP2 Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">Q6P179</a>
Other Accession	<a href="#">NP_071745</a> , <a href="#">11641261</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	Predicted: 106 kDa
Application Notes	Observed: 120kDa KDa ERAP2 antibody can be used for detection of ERAP2 by Western blot at 1 - 2 µg/ml.

**ERAP2 Antibody - Additional Information**Gene ID **64167****Target/Specificity**

ERAP2; ERAP2 antibody is human reactive. Multiple isoforms of ERAP2 are known to exist. ERAP2 antibody is predicted to not cross-react with ERAP1.

**Reconstitution & Storage**

ERAP2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.

**Precautions**

ERAP2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**ERAP2 Antibody - Protein Information****Name** ERAP2**Synonyms** LRAP**Function**

Aminopeptidase that plays a central role in peptide trimming, a step required for the generation of most HLA class I-binding peptides. Peptide trimming is essential to customize longer precursor peptides to fit them to the correct length required for presentation on MHC class I molecules. Preferentially hydrolyzes the basic residues Arg and Lys.

**Cellular Location**

Endoplasmic reticulum membrane; Single-pass type II membrane protein

**Tissue Location**


Ubiquitously expressed. Highly expressed in spleen and leukocytes.

## ERAP2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## ERAP2 Antibody - Images



Western blot analysis of ERAP2 in human ovary tissue lysate with ERAP2 antibody at 1 µg/ml.

## ERAP2 Antibody - Background

The endoplasmic reticulum (ER) aminopeptidase 2 (ERAP2), a member of the peptidase M1 family, like the related protein ERAP1, plays a central role in peptide trimming, a step required for the generation of most HLA class I-binding peptides (1,2). Like ERAP1, ERAP2 is localized to the lumen of the ER and is thought to associate with ERAP1 as a heterodimer (1). Both ERAP1 and ERAP2 have been linked to several human diseases ranging from infections to autoimmunity and cancer, and may play a role in the innate immune response (reviewed in 3).

## ERAP2 Antibody - References

Saveanu L, Carroll O, Lindo V, et al. Concerted peptide trimming by ERAP1 and ERAP2 aminopeptidase complexes in the endoplasmic reticulum. *Nat. Immunol.* 2005; 6:689-97.  
Saric T, Chang SC, Hattori A, et al. An IFN-gamma induced aminopeptidase in the ER, ERAP1, trims precursors to MHC class I-presented peptides. *Nat. Immunol.* 2002; 3:1169-76.  
Cifaldi L, Romania P, Lorenzi S, et al. Role of endoplasmic reticulum aminopeptidases in health and disease: from infection to cancer. *Int. J. Mol. Sci.* 2012; 13:8338-52