

THAP11 Antibody
Catalog # ASC11738**Specification****THAP11 Antibody - Product Information**

Application	WB, E
Primary Accession	Q96EK4
Other Accession	NP_065190 , 40354197
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	Predicted: 35 kDa
Application Notes	Observed: 38 kDa KDa THAP11 antibody can be used for detection of THAP11 by Western blot at 1 - 2 µg/ml.

THAP11 Antibody - Additional Information

Gene ID **57215**
Target/Specificity
THAP11; THAP11 antibody is human, mouse and rat reactive.

Reconstitution & Storage

THAP11 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.

Precautions

THAP11 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

THAP11 Antibody - Protein Information

Name THAP11

Function

Transcription factor, which has both transcriptional activation and repression activities (PubMed:31905202). Also modulates chromatin accessibility (PubMed:38361031). In complex with HCFC1 and ZNF143, regulates the expression of several genes, including AP2S1, ESCO2, OPHN1, RBL1, UBXL1 and ZNF32 (PubMed:26416877). May regulate the expression of genes that encode both cytoplasmic and mitochondrial ribosomal proteins (By similarity). Required for normal mitochondrial development and function. Regulates mitochondrial gene expression, including that of components of the electron transport chain (By similarity). Involved in the maintenance of pluripotency in early embryonic cells, possibly through its action on mitochondrial maturation which is required to meet high energy demands of these cells (By similarity). Required for early development of retina, preventing premature exit of retinal progenitor cells from the cell cycle.

This effect may also be mediated by its action on mitochondria (By similarity). Through the regulation of MMACHC gene expression, controls cobalamin metabolism (PubMed:28449119, PubMed:31905202). Required for normal brain development and neural precursor differentiation (By similarity). Involved in cell growth (PubMed:31905202).

Cellular Location

Nucleus. Cytoplasm Note=In oocytes, detected in the ooplasm, without evidence of its presence in the nucleus (By similarity). Found in the nucleus of undifferentiated embryonic stem cells (PubMed:18585351). Evenly distributed between nucleus and cytoplasm in skin fibroblasts (PubMed:37148549). {ECO:0000250|UniProtKB:Q9JJD0, ECO:0000269|PubMed:18585351, ECO:0000269|PubMed:37148549}

Tissue Location


Expressed in skin fibroblasts.

THAP11 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

THAP11 Antibody - Images



Western blot analysis of THAP11 in human brain tissue lysate with THAP11 antibody at 1 µg/ml.

THAP11 Antibody - Background

The THAP domain-containing protein 11 (THAP11), also known as Ronin, is an essential factor

involved in embryonic stem (ES) cell pluripotency and cell growth (1). THAP 11 contains a THAP domain, a conserved DNA-binding domain common to many proteins associated with chromatin modification and gene expression silencing, and has striking similarity to the site-specific DNA-binding domain (DBD) of Drosophila P element transposases (2). THAP11 can also negatively regulate CD44 v6 expression through its interaction with the poly(rC) binding protein PCBP1 (3).

THAP11 Antibody - References

Dejosez M, Krumenacker JS, Zitun LJ, et al. Ronin is essential for embryogenesis and the pluripotency of mouse ES cells. *Cell* 2008; 133:1162-74.
Roussigne M, Kossida S, Lavigne AC, et al. The THAP domain: a novel protein motif with similarity to the DNA-binding domain of P element transposase. *Trends Biochem. Sci.* 2003; 28:66-9.
Lian WX, Yin RH, Kong XZ, et al. THAP11, a novel binding protein of PCBP1, negatively regulates CD44 alternative splicing and cell invasion in a human hepatoma cell line. *FEBS Lett.* 2012; 586:1431-8.