

IFIT1 Antibody
Catalog # ASC11787**Specification**

IFIT1 Antibody - Product Information

Application
Primary Accession
Other Accession
Reactivity
Host
Clonality
Isotype
Calculated MW

WB, IHC, IF
[P09914](#)
[NP_001539](#), [116534937](#)
Human, Mouse, Rat
Rabbit
Polyclonal
IgG
Predicted: 53 kDa

Application Notes

Observed: 53 kDa KDa
IFIT1 antibody can be used for detection of IFIT1 by Western blot at 1 - 2 µg/ml. Antibody can also be used for Immunohistochemistry at 5 µg/mL. For Immunofluorescence start at 20 µg/mL.

IFIT1 Antibody - Additional Information

Gene ID

3434**Target/Specificity**

IFIT1; IFIT1 antibody is human, mouse and rat reactive. At least three isoforms of IFIT1 are known to exist. This antibody is predicted to not cross-react with other members of the IFIT protein family.

Reconstitution & Storage

IFIT1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.

Precautions

IFIT1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

IFIT1 Antibody - Protein Information**Name** IFIT1**Synonyms** G10P1, IFI56, IFNAI1, ISG56**Function**

Interferon-induced antiviral RNA-binding protein that specifically binds single-stranded RNA bearing a 5'-triphosphate group (PPP-RNA), thereby acting as a sensor of viral single-stranded RNAs and inhibiting expression of viral messenger RNAs. Single-stranded PPP- RNAs, which lack 2'-O-methylation of the 5' cap and bear a 5'- triphosphate group instead, are specific from viruses, providing a molecular signature to distinguish between self and non-self mRNAs by the host during viral infection. Directly binds PPP-RNA in a non- sequence-specific manner. Viruses evolved several

ways to evade this restriction system such as encoding their own 2'-O-methylase for their mRNAs or by stealing host cap containing the 2'-O-methylation (cap snatching mechanism). Exhibits antiviral activity against several viruses including human papilloma and hepatitis C viruses.


Cellular Location

Cytoplasm


IFIT1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

IFIT1 Antibody - Images

Western blot analysis of IFIT1 in rat small intestine tissue lysate with IFIT1 antibody at 1 µg/ml.



Immunohistochemistry of IFIT1 in human small intestine tissue with IFIT1 antibody at 5 µg/mL.



Immunofluorescence of IFIT1 in human small intestine tissue with IFIT1 antibody at 20 µg/mL.

IFIT1 Antibody - Background

The interferon-induced protein with tetratricopeptide repeats 1 (IFIT1) protein is a member of a family of tetratricopeptide repeat-containing proteins whose transcription is upregulated by interferons, virus infection, and molecular patterns such as dsRNA or lipopolysaccharides (1,2). These proteins have been suggested to induce anti-viral cellular activities in response to infection (2). Together with the interferon-induced transmembrane protein 1 (IFITM1), IFIT1 inhibits the replication of the Hepatitis C virus, suggesting that it may be useful in the development of therapeutic treatments (3).

IFIT1 Antibody - References

Wathelet M, Moutschen S, Defilippi P, et al. Molecular cloning, full-length sequence and preliminary characterization of a 56-kDa protein induced by human interferons. *Eur. J. Biochem.* 1986; 155:11-7.

Fensterl V and Sen GC. The ISG56/IFIT1 gene family. *J. Interferon Cytokine Res.* 2011; 31:71-8.

Raychoudhuri A, Shrivastava S, Steele R, et al. ISG56 and IFITM1 proteins inhibit hepatitis C virus replication. *J. Virol.* 2011; 85:12881-9.