

Vinculin Antibody
Catalog # ASC11803**Specification****Vinculin Antibody - Product Information**

Application	WB, IF, ICC, E
Primary Accession	P18206
Other Accession	NP_003364 , 7414
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	Predicted: 120 kDa

Application Notes	Observed: 117 kDa KDa Vinculin antibody can be used for detection of Vinculin by Western blot at 0.5 - 1 µg/ml. For immunofluorescence start at 20 µg/mL.
-------------------	--

Vinculin Antibody - Additional Information

Gene ID **7414**

Target/Specificity

Vinculin antibody was raised against a 23 amino acid peptide near the center of human Vinculin. The immunogen is located within amino acids 820 - 870 of Vinculin.

Reconstitution & Storage

Vinculin antibody can be stored at 4°C for three months and -20°C, stable for up to one year.

Precautions

Vinculin Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Vinculin Antibody - Protein Information

Name VCL

Function

Actin filament (F-actin)-binding protein involved in cell- matrix adhesion and cell-cell adhesion. Regulates cell-surface E- cadherin expression and potentiates mechanosensing by the E-cadherin complex. May also play important roles in cell morphology and locomotion.

Cellular Location

Cell membrane {ECO:0000250|UniProtKB:P12003}; Peripheral membrane protein {ECO:0000250|UniProtKB:P12003}; Cytoplasmic side {ECO:0000250|UniProtKB:P12003}. Cell junction, adherens junction {ECO:0000250|UniProtKB:P12003}. Cell junction, focal adhesion {ECO:0000250|UniProtKB:P12003}. Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:P85972}. Cell membrane, sarcolemma {ECO:0000250|UniProtKB:Q64727}; Peripheral membrane protein

{ECO:0000250|UniProtKB:Q64727}; Cytoplasmic side {ECO:0000250|UniProtKB:Q64727}. Cell projection, podosome {ECO:0000250|UniProtKB:Q64727}. Note=Recruitment to cell-cell junctions occurs in a myosin II-dependent manner. Interaction with CTNNB1 is necessary for its localization to the cell-cell junctions {ECO:0000250|UniProtKB:P12003}


Tissue Location

Metavinculin is muscle-specific.


Vinculin Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Vinculin Antibody - Images

Immunofluorescence of RAIDD in HeLa cells with RAIDD antibody at 5 µg/ml.



Immunohistochemistry of LMX1A in mouse brain tissue with LMX1A Antibody at 5 µg/mL.

Vinculin Antibody - Background

Vinculin is a cytoskeletal protein that plays an important role in the regulation of focal adhesions and embryonic development (1). Three structural vinculin domains include an amino-terminal head, a short flexible proline-rich region and a carboxy-terminal tail (2). Expression of vinculin were shown to be affected by the level of actin expression (2,3). Vinculin deficiencies are associated with a decrease in cell adhesion and an increase in cell motility, suggesting a possible role in metastatic growth (4). Defects in VCL are the cause of cardiomyopathy dilated type 1W (CMD1W) (5).

Vinculin Antibody - References

Burridge K, Fath K, Kelly T, et al. Focal adhesions: transmembrane junctions between the extracellular matrix and the cytoskeleton. Annu. Rev. Cell Biol. 1988; 4:487-525.
Gilmore AP, Jackson P, Waites GT, et al. Further characterization of the talin-binding site in the cytoskeletal protein vinculin. J. Cell Sci. 1992; 103:719-31.
Deakin NO, Ballestrem C, and Turner CE. Paxillin and Hic-5 interaction with vinculin is differentially regulated by Rac1 and RhoA. PLoS One 2012; 7:e37990.
Goldmann WH, Auernheimer V, Thievessen I, et al. Vinculin, cell mechanics and tumour cell invasion. Cell Biol. Int. 2013; Feb 1.