

NOX1 Antibody
Catalog # ASC11832**Specification**

NOX1 Antibody - Product Information

Application	WB, IHC-P, IF, E
Primary Accession	Q9Y5S8
Other Accession	NP_008983 , 148536873
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	Predicted: 62 kDa

Application Notes	Observed: 64 kDa KDa NOX1 antibody can be used for detection of NOX1 by Western blot at 1 - 2 µg/ml. Antibody can also be used for Immunohistochemistry starting at 5 µg/mL. For immunofluorescence start at 20 µg/mL.
-------------------	--

NOX1 Antibody - Additional Information

Gene ID	27035
Target/Specificity	
NOX1; NOX1 antibody is human specific.	

Reconstitution & Storage

NOX1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.

Precautions

NOX1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

NOX1 Antibody - Protein Information

Name NOX1 ([HGNC:7889](#))

Synonyms MOX1, NOH1

Function

NADPH oxidase that catalyzes the generation of superoxide from molecular oxygen utilizing NADPH as an electron donor.

Cellular Location

Cell projection, invadopodium membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein

Tissue Location


[Isoform NOH-1L]: Detected in colon, uterus, prostate, and colon carcinoma, but not in peripheral blood leukocytes

NOX1 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


NOX1 Antibody - Images



Western blot analysis of NOX1 in 293 cell lysate with NOX1 antibody at 1 µg/ml.



Immunohistochemistry of NOX1 in human kidney tissue with NOX1 antibody at 5 µg/ml.



Immunofluorescence of NOX1 in human kidney tissue with NOX1 antibody at 20 µg/ml.

NOX1 Antibody - Background

Voltage-gated proton (hydrogen) channels play an important role in cellular defense against acidic stress (1). NOX1 is a homolog of the catalytic subunit of the superoxide-generating NADPH oxidase of phagocytes, gp91phox (1). Three splice variants of NOX1 have been identified, NOH-1L, NOH-1S and NOH-1Lv (2). NOH-1S is a voltage-gated proton channel that participates in the regulation of cellular pH and is blocked by zinc. NOH-1L is a pyridine nucleotide-dependent oxidoreductase that generates superoxide and might conduct H(+) ions as part of its electron transport mechanism, whereas NOH-1S does not contain an electron transport chain (1-3). NOX1 have the potential to be effective treatments for a range of ischemic diseases (4).

NOX1 Antibody - References

Helmcke I, Heumuller S, Tikkanen R, et al. Identification of structural elements in Nox1 and Nox4 controlling localization and activity. *Antioxid. Redox Signal.* 2009; 11:1279-87.
Piccoli C, D'Aprile A, Ripoli M, et al. Bone-marrow derived hematopoietic stem/progenitor cells express multiple isoforms of NADPH oxidase and produce constitutively reactive oxygen species. *Biochem. Biophys. Res. Commun.* 2007; 353:965-72.
Lee JG, Lim EJ, Park DW, et al. A combination of Lox-1 and Nox1 regulates TLR9-mediated foam cell formation. *Cell Signal.* 2008; 20:2266-75.
Stanic B, Katsuyama M, and Miller FJ Jr. An oxidized extracellular oxidation-reduction state increases Nox1 expression and proliferation in vascular smooth muscle cells via epidermal growth factor receptor activation. *Arterioscler. Thromb. Vasc. Biol.* 2010; 30:2234-41.