

**EDA1 Antibody**  
**Catalog # ASC11876****Specification**

---

**EDA1 Antibody - Product Information**

Application	WB, IHC-P, IF, E
Primary Accession	<a href="#">Q92838</a>
Other Accession	<a href="#">NP_001390</a> , <a href="#">4503449</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	Predicted: 16, 43 kDa

## Application Notes

**Observed: 39 kDa KDa**  
**EDA1 antibody can be used for detection of EDA1 by Western blot at 1 - 2 µg/ml.**  
**Antibody can also be used for immunohistochemistry starting at 5 µg/mL.**  
**For immunofluorescence start at 20 µg/mL.**

**EDA1 Antibody - Additional Information**Gene ID **1896****Target/Specificity**

EDA; EDA1 antibody is human, mouse and rat reactive. Multiple isoforms of EDA1 are known to exist.

**Reconstitution & Storage**

EDA1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.

**Precautions**

EDA1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**EDA1 Antibody - Protein Information****Name** EDA**Synonyms** ED1, EDA2**Function**

Cytokine which is involved in epithelial-mesenchymal signaling during morphogenesis of ectodermal organs. Functions as a ligand activating the DEATH-domain containing receptors EDAR and EDA2R (PubMed:<[a href="http://www.uniprot.org/citations/11039935" target="\\_blank">a href="http://www.uniprot.org/citations/11039935" target="\\_blank">11039935](http://www.uniprot.org/citations/11039935)</a>, PubMed:<[a href="http://www.uniprot.org/citations/27144394" target="\\_blank">a href="http://www.uniprot.org/citations/27144394" target="\\_blank">27144394](http://www.uniprot.org/citations/27144394)</a>, PubMed:<[a href="http://www.uniprot.org/citations/34582123" target="\\_blank">a href="http://www.uniprot.org/citations/34582123" target="\\_blank">34582123](http://www.uniprot.org/citations/34582123)</a>, PubMed:<[a href="http://www.uniprot.org/citations/8696334" target="\\_blank">a href="http://www.uniprot.org/citations/8696334" target="\\_blank">8696334](http://www.uniprot.org/citations/8696334)</a>). May also play a role in cell adhesion (By similarity).

**Cellular Location**

Cell membrane {ECO:0000250|UniProtKB:O54693}; Single-pass type II membrane protein {ECO:0000250|UniProtKB:O54693}


**Tissue Location**

Not abundant; expressed in specific cell types of ectodermal (but not mesodermal) origin of keratinocytes, hair follicles, sweat glands. Also in adult heart, liver, muscle, pancreas, prostate, fetal liver, uterus, small intestine and umbilical cord {ECO:0000269|Ref.6}


**EDA1 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**EDA1 Antibody - Images**

Western blot analysis of EDA1 in mouse kidney tissue lysate with EDA1 antibody at 1  $\mu$ g/ml in (A) the absence and (B) the presence of blocking peptide.



Immunohistochemistry of EDA1 in mouse kidney tissue with EDA1 antibody at 5 µg/ml.



Immunofluorescence of EDA1 in mouse kidney tissue with EDA1 antibody at 20 µg/ml.

### **EDA1 Antibody - Background**

Ectodysplasin A (EDA1) is a member of the TNF-related ligand family involved in the early epithelial-mesenchymal interaction that regulates ectodermal appendage formation (1). It is a trimeric type II membrane protein that co-localizes with cytoskeletal structures at the lateral and apical surfaces of cells and can be expressed as eight alternatively spliced isoforms in hair follicles and in the epidermis of adult skin (2,3). EDAs are required during development, and loss or mutation of EDA1 results in a group of developmental disorders identified as ectodermal dysplasia type 1 (4,5).

### **EDA1 Antibody - References**

Kere J, Srivastava AK, Montonen O. X-linked anhidrotic (hypohidrotic) ectodermal dysplasia is caused by mutation in a novel transmembrane protein. *Nat. Genet.* 1996; 13:409-16.  
Vincent MC, Biancalana V, Ginisty D, et al. Mutational spectrum of the ED1 gene in X-linked hypohidrotic ectodermal dysplasia. *Eur. J. Hum. Genet.* 2001; 9:355-63.  
Ohashi M, Moriya C, Tanahashi K, et al. A new EDA gene mutation in a family of X-linked hypohidrotic ectodermal dysplasia. *J. Dermatol. Sci.* 2014; 74:175-7.  
Bayés M, Hartung AJ, Ezer S, et al. The anhidrotic ectodermal dysplasia gene (EDA) undergoes alternative splicing and encodes ectodysplasin-A with deletion mutations in collagenous repeats. *Hum. Mol. Genet.* 1998; 7:1661-9.