

DTX4 Antibody
Catalog # ASC11879**Specification**

DTX4 Antibody - Product Information

Application	WB, IHC-P, IF, E
Primary Accession	Q9Y2E6
Other Accession	NP_055992 , 148237498
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	Predicted: 68 kDa

Application Notes	Observed: 68 kDa KDa DTX4 antibody can be used for detection of DTX4 by Western blot at 1 - 2 µg/ml. Antibody can also be used for immunohistochemistry starting at 5 µg/mL. For immunofluorescence start at 20 µg/mL.
-------------------	--

DTX4 Antibody - Additional Information

Gene ID **23220**

Target/Specificity

DTX4; DTX4 antibody is human and mouse reactive. At least two isoforms of DTX4 are known to exist; this antibody will detect both isoforms. DTX4 antibody is predicted to not cross-react with other DTX protein family members.

Reconstitution & Storage

DTX4 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.

Precautions

DTX4 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

DTX4 Antibody - Protein Information

Name DTX4

Synonyms KIAA0937, RNF155

Function

Regulator of Notch signaling, a signaling pathway involved in cell-cell communications that regulates a broad spectrum of cell-fate determinations (By similarity). Functions as a ubiquitin ligase protein in vivo, mediating 'Lys48'-linked polyubiquitination and promoting degradation of TBK1, targeting to TBK1 requires interaction with NLRP4.

Cellular Location


Cytoplasm.

DTX4 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


DTX4 Antibody - Images



Western blot analysis of DTX4 in HeLa cell lysate with DTX4 antibody at 1 µg/ml.



Immunohistochemistry of DTX4 in human spleen tissue with DTX4 antibody at 5 µg/ml.



Immunofluorescence of DTX4 in human spleen tissue with DTX4 antibody at 20 µg/ml.

DTX4 Antibody - Background

The Deltex 4 E3 ubiquitin ligase (DTX4) is a member of a protein family that is involved in Notch signaling and neurogenesis (1). Recent studies have shown that DTX4 is also associated with the innate immune response (2). Following the activation of type I interferon signaling, the Nod-like receptor protein 4 (NLRP4) recruits DTX4 for Lys48-linked polyubiquitination and degradation of the kinase TBK1 (NAK), thereby maintaining immune homeostasis during antiviral innate immunity (2).

DTX4 Antibody - References

Kishi N, Tang Z, Maeda Y, et al. Murine homologs of deltex define a novel gene family involved in vertebrate Notch signaling and neurogenesis. *Int. J. Dev. Neurosci.* 2001; 19:21-35.
Cui J, Li Y, Zhu L, et al. NLRP4 negatively regulated type I interferon signaling by targeting the kinase TBK1 for degradation via the ubiquitin ligase DTX4. *Nat. Immunol.* 2012; 13:387-95.