

ATG4B Antibody Catalog # ASC11884

Specification

ATG4B Antibody - Product Information

Application	WB, IHC-P, IF, E
Primary Accession	Q9Y4P1
Other Accession	NP_037457 , 47132611
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	Predicted: 43 kDa
Application Notes	<p>Observed: 44 kDa</p> <p>ATG4B antibody can be used for detection of ATG4B by Western blot at 1 - 2 µg/ml. Antibody can also be used for immunohistochemistry starting at 5 µg/mL. For immunofluorescence start at 20 µg/mL.</p>

ATG4B Antibody - Additional Information

Gene ID	23192
Target/Specificity	ATG4B; ATG4B antibody is human, mouse and rat reactive. At least two isoforms of ATG4B are known to exist; this antibody will detect only the larger isoform. ATG4B is predicted to not cross-react with other ATG4 proteins.

Reconstitution & Storage

ATG4B antibody can be stored at 4°C for three months and -20°C, stable for up to one year.

Precautions

ATG4B Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

ATG4B Antibody - Protein Information

Name ATG4B {ECO:0000303|PubMed:15187094, ECO:0000312|HGNC:HGNC:20790}

Function

Required for canonical autophagy (macroautophagy), non-canonical autophagy as well as for mitophagy (PubMed:33773106, PubMed:33909989). The protease activity is required for proteolytic activation of ATG8 family proteins: cleaves the C-terminal amino acid of ATG8 proteins MAP1LC3A, MAP1LC3B, MAP1LC3C, GABARAPL1, GABARAPL2 and GABARAP, to reveal a C-terminal glycine (PubMed:15169837, PubMed:15187094, PubMed:17347651, PubMed:19322194, PubMed:20818167, PubMed:21177865, PubMed:22302004, PubMed:27527864, PubMed:28287329, PubMed:28633005, PubMed:29458288, PubMed:30661429). Exposure of the glycine at the C-terminus is essential for ATG8 proteins conjugation to phosphatidylethanolamine (PE) and insertion to membranes, which is necessary for autophagy (PubMed:15169837, PubMed:15187094, PubMed:17347651, PubMed:19322194, PubMed:21177865, PubMed:22302004). Protease activity is also required to counteract formation of high-molecular weight conjugates of ATG8 proteins (ATG8ylation): acts as a deubiquitinating-like enzyme that removes ATG8 conjugated to other proteins, such as ATG3 (PubMed:31315929, PubMed:33773106). In addition to the protease activity, also mediates delipidation of ATG8 family proteins (PubMed:15187094, PubMed:19322194, PubMed:28633005, PubMed:29458288, PubMed:32686895, PubMed:33909989). Catalyzes delipidation of PE-conjugated forms of ATG8 proteins during macroautophagy (PubMed:15187094, PubMed:19322194, PubMed:29458288, PubMed:32686895, PubMed:33909989). Also involved in non-canonical autophagy, a parallel pathway involving conjugation of ATG8 proteins to single membranes at endolysosomal compartments, by catalyzing delipidation of ATG8 proteins conjugated to phosphatidylserine (PS) (PubMed:33909989). Compared to other members of the family (ATG4A, ATG4C or ATG4C), constitutes the major protein for

proteolytic activation of ATG8 proteins, while it displays weaker delipidation activity than other ATG4 paralogs (PubMed:29458288, PubMed:30661429). Involved in phagophore growth during mitophagy independently of its protease activity and of ATG8 proteins: acts by regulating ATG9A trafficking to mitochondria and promoting phagophore-endoplasmic reticulum contacts during the lipid transfer phase of mitophagy (PubMed:33773106).

Cellular Location


Cytoplasm. Cytoplasm, cytosol. Cytoplasmic vesicle, autophagosome. Endoplasmic reticulum. Mitochondrion. Note=Mainly localizes to the cytoplasm, including cytosol (PubMed:29165041). A small portion localizes to mitochondria; phosphorylation at Ser-34 promotes localization to mitochondria (PubMed:29165041).

ATG4B Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


ATG4B Antibody - Images



Western blot analysis of ATG4B in 3T3 cell lysate with ATG4B antibody at 1 µg/ml.



Immunohistochemistry of ATG4B in human spleen tissue with ATG4B antibody at 5 µg/ml.



Immunofluorescence of ATG4B in human spleen tissue with ATG4B antibody at 20 µg/ml.

ATG4B Antibody - Background

Autophagy, the process of bulk degradation of cellular proteins through an autophagosomes-lysosomal pathway is important for normal growth control and may be defective in tumor cells. It is involved in the preservation of cellular nutrients under starvation conditions as well as the normal turnover of cytosolic components (1,2). ATG4B, also known as AUTL1, is one of four mammalian orthologs of the yeast ATG4 protein; all four are cysteine proteases (3). ATG4 is required for ATG8 conjugation to phosphatidylethanolamine on autophagosomal membranes. In mammals, each ATG4 homolog shows a selective preference for the ATG8 homologs (4). ATG4B has been found to be a novel protective protein in inflammatory colitis (5).

ATG4B Antibody - References

- Gozuacik D and Kimchi A. Autophagy as a cell death and tumor suppressor mechanism. *Oncogene* 2004; 23:2891-906.
- Kisen GO, Tessitore L, Costelli P, et al. Reduced autophagic activity in primary rat hepatocellular carcinoma and ascites hepatoma cells. *Carcinogenesis* 1993; 14:2501-5.
- Marino G, Uria JA, Puente XS, et al. Human autophagins, a family of cysteine proteinases potentially implicated in cell degradation by autophagy. *J. Biol. Chem.* 2003; 278:3671-8.
- Li M, Hou Y, Wang J, et al. Kinetic comparisons of mammalian Atg4 homologues indicate selective preferences towards diverse Atg8 substrates. *J. Biol. Chem.* 2011; 286:7327-38.