

# **DNAJC13** Antibody

Catalog # ASC11897

# Specification

# **DNAJC13 Antibody - Product Information**

Application **Primary Accession** Other Accession Reactivity Host Clonality Isotype Calculated MW

**Application Notes** 

WB, IHC 075165 NP 056083, 112421122 Human, Mouse Rabbit Polyclonal laG Predicted: 249 kDa

**Observed: 249 kDa KDa** DNAJC13 antibody can be used for detection of DNAJC13 by Western blot at 1 - 2 µg/mL. Antibody can also be used for immunohistochemistry starting at 5 µg/mL.

### **DNAJC13 Antibody - Additional Information**

Gene ID

**Target/Specificity** 

DNAJC13; DNAJC13 antibody is human and mouse reactive. At least two siforms of DNAJC13 are known to exist; this antibody will detect both isoforms. DNAJC13 antibody is predicted to not cross-react with other DNAJC family members.

23317

#### **Reconstitution & Storage**

DNAJC13 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.

**Precautions** DNAJC13 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

# **DNAJC13 Antibody - Protein Information**

Name DNAJC13

Synonyms KIAA0678, RME8

#### **Function**

Involved in membrane trafficking through early endosomes, such as the early endosome to recycling endosome transport implicated in the recycling of transferrin and the early endosome to late endosome transport implicated in degradation of EGF and EGFR (PubMed: <a href="http://www.uniprot.org/citations/18256511" target="\_blank">18256511</a>, PubMed:<a href="http://www.uniprot.org/citations/18307993" target="\_blank">18200011 </d> the regulation of endosomal membrane tubulation and regulates the dynamics of SNX1 on the endosomal membrane; via association with WASHC2 may link the WASH complex to the retromer



SNX-BAR subcomplex (PubMed:<a href="http://www.uniprot.org/citations/24643499" target="\_blank">24643499</a>).

# **Cellular Location**

Early endosome. Early endosome membrane; Peripheral membrane protein. Endosome membrane

# **DNAJC13 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- <u>Flow Cytomety</u>
- <u>Cell Culture</u>

# **DNAJC13** Antibody - Images



Western blot analysis of DNAJC13 in human heart tissue lysate with DNAJC13 antibody at 1 µg/ml.





Immunohistochemistry of DNAJC13 in human heart tissue with DNAJC13 antibody at 5  $\mu$ g/mL.

# DNAJC13 Antibody - Background

DNAJC13, also known as receptor-mediated endocytosis 8 (RME8), is the human homolog to a DnaJ domain-containing protein originally identified in a screen for endocytic defects in C. elegans (1). It is thought to be a co-chaperone of Hsc70 which regulates protein conformation at membrane sites and plays a role in intracellular trafficking, co-localizing with markers of the endosomal system. Recent experiments have indicated that the DNAJC13 protein is involved in membrane trafficking through early endosomes but not through degradative organelles (2). DNAJC13 has been also been shown to regulate the intracellular trafficking of the epidermal growth factor receptor (3).

#### **DNAJC13 Antibody - References**

Girard M, Poupon V, Blondeau F, et al. The DnaJ-domain protein RME-8 functions in endosomal trafficking. J. Biol. Chem. 2005; 280:40135-43.

Fujibayashi A, Taguchi T, Misaki R, et al. Human RME-8 is involved in membrane trafficking through early endosomes. Cell Struct. Funct. 2008; 33:35-50.

Girard M and McPherson PS. RME-8 regulates trafficking of the epidermal growth factor receptor. FEBS Lett. 2008; 582:961-6.