

BHLHA15 Antibody

Catalog # ASC12032

Specification

BHLHA15 Antibody - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Isotype Calculated MW Application Notes WB <u>Q7RTS1</u> <u>29126247, NP_803238, 168620</u> Human, Mouse, Rat Rabbit Polyclonal IgG 20818 BHLHA15 antibody can be used for Western blot at 1 - 2 µg/mL.

BHLHA15 Antibody - Additional Information

Gene ID 168620 Other Names BHLHA15 Antibody: FB22, HM89, LAP3, LCR1, NPYR, WHIM, CD184, LESTR, NPY3R, NPYRL, HSY3RR, NPYY3R, D2S201E

Precautions BHLHA15 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

BHLHA15 Antibody - Protein Information

Name BHLHA15

Synonyms BHLHB8, MIST1

Function

Plays a role in controlling the transcriptional activity of MYOD1, ensuring that expanding myoblast populations remain undifferentiated. Repression may occur through muscle-specific E-box occupancy by homodimers. May also negatively regulate bHLH-mediated transcription through an N-terminal repressor domain. Serves as a key regulator of acinar cell function, stability, and identity. Also required for normal organelle localization in exocrine cells and for mitochondrial calcium ion transport. May function as a unique regulator of gene expression in several different embryonic and postnatal cell lineages. Binds to the E-box consensus sequence 5'-CANNTG-3' (By similarity).

Cellular Location Nucleus.

Tissue Location Expressed in brain, liver, spleen and skeletal muscle.




BHLHA15 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

BHLHA15 Antibody - Images



Western blot analysis of BHLHA15 in 3T3 cell lysate with BHLHA15 antibody at (A) 1 and (B) 2 $\mu g/mL$

BHLHA15 Antibody - Background

BHLHA15 Antibody: BHLHA15 (basic helix-loop-helix protein 15), also known as MIST1 (muscle intestine and stomach expression 1) belongs to the bHLH family of transcription factors and plays a role in regulating the transcriptional activity of MYOD1 in muscle cell development as well as serving as a key regulator of acinar cell function (1,2). BHLHA15 contains a basic helix-loop-helix (bHLH) domain and is capable of binding to E-box motifs as a homodimer or a heterodimer with E-proteins. It may also negatively regulate bHLH-mediated transcription through a N-terminal repressor domain (3). It is expressed in mammary epithelial cells and is essential for the regulation of mammary gland development (4).

BHLHA15 Antibody - References

Lemercier C, To RQ, Swanson BJ et al. MIST1: a novel basic helix-loop-helix transcription factor exhibits a developmentally regulated expression pattern. Dev. Biol. 1997; 182:101-13.;Pin CL, Rukstalis JM, Johnson C, et al. The bHLH transcription factor MIST1 is required to maintain exocrine pancreas cell organization and acinar cell identity. J. Cell Biol. 2001; 155:519-30.;Lemercier C, To RQ, Carrasco RA et al. The basic helix-loop-helix transcription factor Mist1 functions as a transcriptional repressor of myoD. EMBO J. 1998; 17:1412-22.;Zhao Y, Johansson C, Tran T, et al.



Identification of a basic helix-loop-helix transcription factor expressed in mammary gland alveolar cells and required for maintenance of the differentiated state. Mol. Endocrinol. 2006; 20:2187-98.