

## **NDST1 Antibody**

Catalog # ASC12107

### **Specification**

## **NDST1 Antibody - Product Information**

**Application Primary Accession** Other Accession Host Clonality Isotype

Calculated MW

WB. E P52848 NP 001534 Rabbit **Polyclonal** 

IaG

Predicted: 97 kDa

Observed: 97 kDa KDa

### **NDST1** Antibody - Additional Information

Gene ID 15531 Alias Symbol NDST1

**Other Names** 

NDST1 Antibody: HSST, NST1, MRT46

# Target/Specificity

NDST1 Antibody is predicted to not cross-react with NDST2.

## **Reconstitution & Storage**

NDST1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

## **Precautions**

NDST1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## **NDST1 Antibody - Protein Information**

Name NDST1 (HGNC:7680)

Synonyms HSST, HSST1

### **Function**

[Isoform 1]: Essential bifunctional enzyme that catalyzes both the N-deacetylation and the N-sulfation of glucosamine (GlcNAc) of the glycosaminoglycan in heparan sulfate (PubMed: <a href="http://www.uniprot.org/citations/35137078" target=" blank">35137078</a>, PubMed:<a href="http://www.uniprot.org/citations/9230113" target=" blank">9230113</a>, PubMed:<a href="http://www.uniprot.org/citations/9744796" target="blank">9744796</a>). Modifies the GlcNAc-GlcA disaccharide repeating sugar backbone to make N-sulfated heparosan, a prerequisite substrate for later modifications in heparin biosynthesis (PubMed: <a

href="http://www.uniprot.org/citations/9230113" target=" blank">9230113</a>). Plays a role in



determining the extent and pattern of sulfation of heparan sulfate. Participates in biosynthesis of heparan sulfate that can ultimately serve as L-selectin ligands, thereby playing a role in inflammatory response (By similarity). Required for the exosomal release of SDCBP, CD63 and syndecan (PubMed:<a href="http://www.uniprot.org/citations/22660413" target="blank">22660413</a>).

#### **Cellular Location**

[Isoform 1]: Golgi apparatus, trans-Golgi network membrane; Single-pass type II membrane protein. Golgi apparatus, cis-Golgi network membrane; Single-pass type II membrane protein

#### **Tissue Location**

Widely expressed. Expression is most abundant in heart, liver and pancreas.

## **NDST1 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## **NDST1** Antibody - Images

## **NDST1 Antibody - Background**

NDST1 is a member of the heparan sulfate/heparin GlcNAc N-deacetylase/ N-sulfotransferase family. It is a type II transmembrane protein that resides in the Golgi apparatus. The encoded protein catalyzes the transfer of sulfate from 3'-phosphoadenosine 5'-phosphosulfate to nitrogen of glucosamine in heparan sulfate.