

VISTA Antibody
Catalog # ASC12122**Specification****VISTA Antibody - Product Information**

Application	WB, IHC, IF, ICC, E
Primary Accession	Q9H7M9
Other Accession	NP_071436
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG

VISTA Antibody - Additional Information

Gene ID	64115
Alias Symbol	VSIR

Other Names

VISTA Antibody: VISTA molecule, VSIR, B7-H5, B7H5, GI24, PP2135, SISP1, DD1alpha, VISTA, C10orf54, chromosome 10 open reading frame 54, PD-1H, V-set immunoregulatory receptor, V-Type Immunoglobulin Domain-Containing Suppressor Of T-Cell Activation, Chromosome 10 Open Reading Frame 54

Reconstitution & Storage

VISTA antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

VISTA Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

VISTA Antibody - Protein Information

Name VSIR ([HGNC:30085](#))

Function

Immunoregulatory receptor which inhibits the T-cell response (PubMed:24691993). May promote differentiation of embryonic stem cells, by inhibiting BMP4 signaling (By similarity). May stimulate MMP14- mediated MMP2 activation (PubMed:20666777).

Cellular Location

Cell membrane; Single-pass type I membrane protein

Tissue Location

Expressed in spleen. Detected on a number of myeloid cells including CD11b monocytes, CD66b+ neutrophils, at low levels on CD4+ and CD8+ T-cells, and in a subset of NK cells. Not detected on B cells (at protein level). Expressed at high levels in placenta, spleen, plasma blood leukocytes,

and lung. Expressed at moderate levels in lymph node, bone marrow, fat, uterus, and trachea Has low expression levels in other tissues

VISTA Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

VISTA Antibody - Images

VISTA Antibody - Background

The lymphocyte activation gene-3 (LAG3) is a member of the immunoglobulin superfamily and binds MHC class II with high affinity (1), negatively regulating T-cell function and homeostasis (2). It is expressed in B, T, and NK cells, monocytes, and dendritic cells (3), and acts to regulate T cell expansion (4). LAG3 is also an important immune checkpoint protein, with anti-LAG3 antibodies activating T effector cells and affecting regulatory T cell functions. Furthermore LAG3 appears to act in a synergistic fashion with PD-1/PD-L1, suggesting that a dual antibody approach may prove useful in cancer immunotherapy (5).

VISTA Antibody - References

Mayya V., et al . Quantitative phosphoproteomic analysis of T cell receptor signaling reveals system-wide modulation of protein-protein interactions. 2009, Sci. Signal. 2:RA46-RA46. Sakr M.A., et al., Gl24 enhances tumor invasiveness by regulating cell surface membrane-type 1 matrix metalloproteinase. 2010, Cancer Sci. 101:2368-2374.