

**CaMKII Antibody**  
**CaMKII Antibody, Clone 22B1**  
**Catalog # ASM10038****Specification****CaMKII Antibody - Product Information**

Application	WB, IHC, ICC, IP, E
Primary Accession	<a href="#">P11275</a>
Other Accession	<a href="#">NP_037052.1</a>
Host	Mouse
Isotype	IgG1
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal

**Description**

Mouse Anti-Rat CaMKII Monoclonal IgG1

**Target/Specificity**Detects phosphorylated CaMKII from rat tissues. This antibody is specific for  $\alpha$  and  $\beta$  subunits of CaMKII only when they are phosphorylated at Thr-286/287 (in  $\beta$ ).**Other Names**

CSAID Binding protein 1 Antibody, CSBP1 Antibody, CSBP2 Antibody, EXIP Antibody, MAP kinase MXI2 Antibody, MAPkinase p38alpha Antibody, MAPK14 Antibody, p38 ALPHA Antibody, p38 MAP kinase Antibody, p38 mitogen activated protein kinase Antibody, RK Antibody, SAPK 2A Antibody, Stress activated protein kinase 2A Antibody

**Immunogen**

Synthetic peptide

**Purification**

Protein G Purified

**Storage****-20°C****Storage Buffer**

PBS pH7.4, 50% glycerol, 0.09% sodium azide

**Shipping Temperature****Blue Ice or 4°C****Certificate of Analysis**1  $\mu$ g/ml was sufficient for detection of 0.2  $\mu$ g CamKII by colorimetric immunoblot analysis using Goat Anti-Mouse IgG:HRP as the secondary.**Cellular Localization**


Cytoplasm | Mitochondrion | Nucleus | Cell Junction | Synapse | Presynaptic Cell Membrane

**CaMKII Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)


- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**CaMKII Antibody - Images**


Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-CaMKII Monoclonal Antibody, Clone 22B1 (ASM10038). Tissue: dissociated hippocampal neurons. Species: Rat. Fixation: Cold 4% paraformaldehyde/0.2% glutaraldehyde in 0.1M sodium phosphate buffer. Primary Antibody: Mouse Anti-CaMKII Monoclonal Antibody (ASM10038) at 1:1000 for 12 hours at 4°C. Secondary Antibody: FITC Goat Anti-Mouse IgG (green) at 1:50 for 30 minutes at RT. Magnification: 10X. Courtesy of: Mary Kennedy, Caltech.



Immunohistochemistry analysis using Mouse Anti-CaMKII Monoclonal Antibody, Clone 22B1 (ASM10038). Tissue: backskin. Species: Mouse. Fixation: Bouin's Fixative and paraffin-embedded. Primary Antibody: Mouse Anti-CaMKII Monoclonal Antibody (ASM10038) at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT. Localization: Muscle, hair follicle, epidermis.



Immunohistochemistry analysis using Mouse Anti-CaMKII Monoclonal Antibody, Clone 22B1 (ASM10038). Tissue: colon carcinoma. Species: Human. Fixation: Formalin. Primary Antibody: Mouse Anti-CaMKII Monoclonal Antibody (ASM10038) at 1:5000 for 12 hours at 4°C. Secondary Antibody: Biotin Goat Anti-Mouse at 1:2000 for 1 hour at RT. Counterstain: Mayer Hematoxylin (purple/blue) nuclear stain at 200  $\mu$ l for 2 minutes at RT. Magnification: 40x.



Western Blot analysis of Mouse Ventrite lysates showing detection of CaMKII protein using Mouse Anti-CaMKII Monoclonal Antibody, Clone 22B1 (ASM10038). Primary Antibody: Mouse Anti-CaMKII Monoclonal Antibody (ASM10038) at 1:1000. Analysis of CaMKII and NFAT phosphorylation in ventricles of 14 day old mice over-expressing CaMK.

#### CaMKII Antibody - Background

CaMKII is an important member of the calcium/calmodulin-activated protein kinase family, functioning in neural synaptic stimulation and T-cell receptor signaling (1, 2). CaMKII is expressed in many different tissues but is specifically found in the neurons of the forebrain and its mRNA is found within the dendrites and the soma of the neuron. The CaMKII that is found in the neurons consist of two subunits of 52 (termed alpha genes) and 60 kDa (beta genes). CaMKII has catalytic and regulatory domains, as well as an ATP-binding domain, and a consensus phosphorylation site (3-7). The binding of Ca<sup>2+</sup>/calmodulin to its regulatory domain releases its auto inhibitory effect and activates the kinase (8). This kinase activation results in autophosphorylation at threonine 286 (8). The threonine phosphorylation state of CaMKII can be regulated through PP1/PKA. Whereas PP1 (protein phosphatase 1) dephosphorylates phospho-CaMKII at Thr286, PKA (protein kinase A) prevents this dephosphorylation (9). Autophosphorylation also enables CaMKII to attain an enhanced affinity for NMDA receptors in postsynaptic densities (10-12).

### CaMKII Antibody - References

1. Hughes K. et al. (2001) *J. Biol. Chem.* 276: 36008-36013.
2. Barria A. et al. (1997) *Science* 276: 2042-2045.
3. Bennet M.K. and Kennedy M.B. (1987) *Proc. Natl. Acad. Sci. U.S.A.* 84: 1794-1798.
4. Broke L., Srinivasan M. and Schulman H. (1995) *J. Neurosci.* 15: 6797-6808.
5. Nghiem P., Saati S. M., Martens C. L., Gardner P. and Schulman H. (1993) *J. Biol. Chem.* 268: 5471-5479.
6. Edman C.F. and Schulman H. (1994) *Biochem. Biophys. Acta* 1221: 90-102.
7. Tombes R.M. and Krystal G.W., (1997) *Biochem. Biophys. Acta* 13555: 281-292.
8. Means A.R. (2000) *Mol. Endocrinol.* 14: 4-12.
9. Makhinson M. et al. (1999) *J. Neurosci.* 19: 2500-2510.
10. Strack S. and Colbran R.J. (1998) *J. Biol. Chem.* 273: 20689-20692.
11. Leonard S.A., Lim I.A., Hemsworth D.E., Horne M.C. and Hell J.W. (1999) *Proc. Natl. Acad. Sci. U.S.A.* 96: 3239-3244.
12. Shen K. and Meyer Y. (1999) *Science* 284: 162-167.