

HSP40, YDJ1 Antibody
HSP40, YDJ1 Antibody, Clone 2A7.H6
Catalog # ASM10099**Specification****HSP40, YDJ1 Antibody - Product Information**

Application	WB, E, IP
Primary Accession	P25491
Other Accession	NP_014335.1
Host	Mouse
Isotype	IgG1 Kappa
Reactivity	Yeast
Clonality	Monoclonal
Format	Biotin

Description

Mouse Anti-Yeast HSP40, YDJ1 Monoclonal IgG1 Kappa

Target/Specificity

Detects ~40kDa. Yeast specific. Does not cross react with Human, Mouse or Rat.

Other Names

Dnaj (HSP40) homolog subfamily B member 1 antibody, DNAJ 1 antibody, DNAJ B1 antibody, Dnaj homolog subfamily B member 1 antibody, Dnaj protein homolog 1 antibody, DNAJ1 antibody, DNAJB 1 antibody, DNAJB1 antibody, DNAJB1 protein antibody, DNJB1_HUMAN antibody, HDJ 1 antibody, HDJ-1 antibody, HDJ1 antibody, Heat shock 40 kDa protein 1 antibody, Heat shock 40kD protein 1 antibody, Heat shock protein 40 antibody, HSP 40 antibody, HSP40 antibody, HSPF 1 antibody, HSPF1 antibody, Human Dnaj protein 1 antibody, Radial spoke 16 homolog B antibody, RSPH16B antibody, Sis1 antibody

Immunogen

Full length protein HSP40 (YDJ1)

Purification

Protein G Purified

Storage

-20°C

Storage Buffer

50% glycerol, 0.09% sodium azide

Shipping Temperature

Blue Ice or 4°C

Certificate of Analysis

0.5 µg/ml of SMC-166 was sufficient for detection of 50 ng YDJ1 by colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody.

Cellular Localization

Cytoplasm | Nucleus

HSP40, YDJ1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HSP40, YDJ1 Antibody - Images



HSP40, YDJ1 Antibody - Background

Human HSP40/Dnaj proteins comprise a large protein family, members of which feature the J domain (named after the bacterial Dnaj protein) (1). The J-domain spans the first 75 N-terminal amino acids and is separated from the C-terminal by a glycine/phenylalanine-rich domain (2). Members of the HSP40/Dnaj family play diverse roles in many cellular processes, such as folding, translocation, degradation and assembly of multi-protein complexes. In particular, Hdj1, the first human HSP40/Dnaj protein identified, plays an important role in protein translation and folding, as well as in the regulation of HSP70 function (3). HSP40 stimulates the ATPase activity of HSP70 which in turn causes conformational changes of the unfolded proteins (4, 5). The HSP40-HSP70-unfolded protein complex further binds to co-chaperones Hip, Hop and HSP90 which leads to protein folding, or components of protein degradation machinery CHIP and BAG-1 (6). Some studies have shown that the difference between HDJ1 and type 1 DNAJ proteins including HDJ2 and yeast Ydj1 is the result of the possession of a zinc finger domain by the latter, which helps in the function of protein folding. (7, 8).

HSP40, YDJ1 Antibody - References

1. Cheetham M.E. and Caplan A.J. (1998) Cell Stress Chaperones 3: 28-36.
2. Fan C.Y., et al. (2003) Cell Stress Chaperones 8: 309-316.
3. Sohn S.Y., Kim S.B., Kim J., and Ahn B.Y. (2006) J Gen Virol. 87(7): 1883-91.
4. Liberek K. et al. (1991) Proc. Natl. Acad. Sci. USA 88: 2874-2878.

5. Cyr D.M., et al. (1992) *J Biol Chem.* 267: 20927-20931.
6. Höhfeld J., et al. (2001) *EMBO Rep.* 2: 885-890.
7. Terda K., et al. (1997) *J Cell Biol.* 139: 1089-1095.
8. Lu Z. and Cyr D.M. (1998) *J Biol Chem.* 273: 27824-27830.