

ADAM22 (extracellular) Antibody
ADAM22 Antibody, Clone S57-2
Catalog # ASM10246**Specification**

ADAM22 (extracellular) Antibody - Product Information

Application	WB, ICC, IP
Primary Accession	O9R1V6
Other Accession	NP_001007221.1
Host	Mouse
Isotype	IgG1
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal

Description

Mouse Anti-Mouse ADAM22 (extracellular) Monoclonal IgG1

Target/Specificity

Detects the extracellular domain of ADAM22 ~90kDa. Does not cross-react with ADAM11. Weakly reactive in human samples.

Other Names

MDC2 Antibody, Disintegrin and metalloproteinase domain-containing protein 22 Antibody, Metalloproteinase disintegrin ADAM22-3 Antibody, Metalloproteinase-like disintegrin-like and cysteine-rich protein 2 Antibody, ADAM 22 Antibody, ADAM metalloproteinase domain 22 Antibody, MGC149832 Antibody

Immunogen

Fusion protein amino acids 444-526 (extracellular disintegrin domain) of mouse ADAM22

Purification

Protein G Purified

Storage **-20°C**

Storage Buffer

PBS pH7.4, 50% glycerol, 0.09% sodium azide

Shipping Temperature

Blue Ice or 4°C

Certificate of Analysis

1 µg/ml of SMC-412 was sufficient for detection of ADAM22 in 10 µg of rat brain lysate by colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody.

Cellular Localization

Membrane


ADAM22 (extracellular) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ADAM22 (extracellular) Antibody - Images



Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-ADAM22 (extracellular) Monoclonal Antibody, Clone S57-2 (ASM10246). Tissue: Neuroblastoma cell line (SK-N-BE). Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Mouse Anti-ADAM22 (extracellular) Monoclonal Antibody (ASM10246) at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:100 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain at 1:1000, 1:5000 for 60min RT, 5min RT. Localization: Membrane. Magnification: 60X. (A) DAPI (blue) nuclear stain (B) Phalloidin Texas Red F-Actin stain (C) ADAM22 (extracellular) Antibody (D) Composite.



Western Blot analysis of Rat brain lysates showing detection of ADAM22 protein using Mouse Anti-ADAM22 Monoclonal Antibody, Clone S57-2 (ASM10246). Primary Antibody: Mouse

Anti-ADAM22 Monoclonal Antibody (ASM10246) at 1:250, 1:500, and 1:1000.

ADAM22 (extracellular) Antibody - Background

ADAM 22 belongs to the ADAM gene family which have been shown to bind integrin and therefore may have a part in cell to cell or cell to matrix interactions. ADAM 22 is unique in the fact that it is only observed in the nervous system and predominantly in the brain. ADAM 22 is attached by cytoskeletal scaffolds to the postsynaptic density and is a receptor for LGI1.

ADAM22 (extracellular) Antibody - References

1. Sagane K., Ohya Y., Hasegawa T., Tanaka I. (1998) Biochem J. 334 (Pt1): 93-98.