

**Neurologin 4 Antibody**  
**Neurologin 4 Antibody, Clone S98-7**  
**Catalog # ASM10301****Specification**

---

**Neurologin 4 Antibody - Product Information**

Application	WB, IHC, ICC
Primary Accession	<a href="#">B0F2B4</a>
Other Accession	<a href="#">ABS19580</a>
Host	Mouse
Isotype	IgG1
Reactivity	Mouse
Clonality	Monoclonal

**Description**

Mouse Anti-Mouse Neurologin 4 Monoclonal IgG1

**Target/Specificity**

Detects ~125-130kDa. Does not cross-react with other Neurologins.

**Other Names**

ASPGX2 Antibody, AUTSX2 Antibody, HLNK Antibody, HNLK Antibody, KIAA1260 Antibody, MGC22376 Antibody, NLGN Antibody, NLGN4 Antibody, NLGN4X Antibody, neurologin-4 X-linked Antibody, neurologin X Antibody, neurologin 4 Antibody, X-linked Antibody

**Immunogen**

Fusion protein amino acids 782-945 (intracellular C-terminus) of mouse Neurologin 4. ~45% identity with Neurologin-1, -2 and -3.

**Purification**

Protein G Purified

Storage **-20°C**

**Storage Buffer**

PBS pH 7.4, 50% glycerol, 0.1% sodium azide

Shipping Temperature

**Blue Ice or 4°C**

**Certificate of Analysis**

1 µg/ml of SMC-469 was sufficient for detection of Neurologin 4 in 20 µg of mouse brain lysate by colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody.

**Cellular Localization**

Cell Membrane | Cell Junction


**Neurologin 4 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Neurologin 4 Antibody - Images



Immunohistochemistry analysis using Mouse Anti-Neurologin 4 Monoclonal Antibody, Clone N98/7 (ASM10301). Tissue: Dentate Gyrus. Species: Mouse. Fixation: formaldehyde-fixed paraffin embedded. Primary Antibody: Mouse Anti-Neurologin 4 Monoclonal Antibody (ASM10301) at 1:100. Secondary Antibody: FITC Goat Anti-Mouse (green). Counterstain: DAPI (blue) nuclear stain at 1:1000. Courtesy of: Rachel Reith, NIH/NIMH.



Western Blot analysis of Mouse Brain Membrane showing detection of ~125-130 kDa Neurologin 4 protein using Mouse Anti-Neurologin 4 Monoclonal Antibody, Clone N98/7 (ASM10301). Lane 1: Molecular Weight Ladder (MW). Lane 2: Mouse Brain Membrane. Load: 15 µg. Block: 2% BSA and 2% Skim Milk in 1X TBST. Primary Antibody: Mouse Anti-Neurologin 4 Monoclonal Antibody (ASM10301) at 1:1000 for 16 hours at 4°C. Secondary Antibody: Goat Anti-Mouse IgG: HRP at 1:2000 for 60 min at RT. Color Development: ECL solution for 6 min at RT. Predicted/Observed Size: ~125-130 kDa.

## Neurologin 4 Antibody - Background

This gene encodes a member of a family of neuronal cell surface proteins. Members of this family may act as splice site-specific ligands for beta-neurexins and may be involved in the formation and remodeling of central nervous system synapses. The encoded protein interacts with discs, large (Drosophila) homolog 4 (DLG4). Mutations in this gene have been associated with autism and Asperger syndrome. Two transcript variants encoding the same protein have been identified for this gene.