

7-Ketocholesterol Antibody
7-Ketocholesterol Antibody, Clone 7E1
Catalog # ASM10340**Specification**

7-Ketocholesterol Antibody - Product Information

Application	ICC/IF, WB
Host	Mouse
Isotype	IgG2b
Clonality	Monoclonal

Description

Mouse Anti-7-Ketocholesterol (7-KC) Monoclonal IgG2b

Target/Specificity

Specific for 7-ketocholesterol modified proteins.

Other Names

7-Ketocholesterol [7-KC] Antibody, 7-KC Antibody, 7-ketocholesterol (7-KC) Antibody, 7KC Antibody, 7KC-Modified Keyhole-Limpet Hemocyanine Antibody, 7KC-Modified KLH Antibody

Trademark

MOLECULAR SIGNATURE®

Immunogen

Synthetic 7-Ketocholesterol conjugated to Keyhole Limpet Hemocyanin (KLH).

Purification

Protein G Purified

Storage

-20°C

Storage Buffer

PBS pH 7.4, 50% glycerol, 0.9% Sodium Azide

Shipping Temperature

Blue Ice or 4°C

Certificate of Analysis


A 1:1000 dilution of SMC-512 was sufficient for detection of 7-Ketocholesterol in 2 µg of 7-ketocholesterol conjugated to BSA by ECL immunoblot analysis using Goat Anti-Mouse IgG:HRP as the secondary Antibody.

7-Ketocholesterol Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

7-Ketocholesterol Antibody - Images



Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-7-Ketocholesterol Monoclonal Antibody, Clone 7E1 (ASM10340). Tissue: Embryonic kidney cells (HEK293). Species: Human. Fixation: 5% Formaldehyde for 5 min. Primary Antibody: Mouse Anti-7-Ketocholesterol Monoclonal Antibody (ASM10340) at 1:50 for 30-60 min at RT. Secondary Antibody: Goat Anti-Mouse Alexa Fluor 488 at 1:1500 for 30-60 min at RT. Counterstain: Phalloidin Alexa Fluor 633 F-Actin stain; DAPI (blue) nuclear stain at 1:250, 1:50000 for 30-60 min at RT. Magnification: 20X (2X Zoom). (A,C,E,G) - Untreated. (B,D,F,H) - Cells cultured overnight with 50 μ M H₂O₂. (A,B) DAPI (blue) nuclear stain. (C,D) Phalloidin Alex Fluor 633 F-Actin stain. (E,F) 7-Ketocholesterol Antibody. (G,H) Composite. Courtesy of: Dr. Robert Burke, University of Victoria.



Western Blot analysis of 7-Ketocholesterol-BSA Conjugate showing detection of 67 kDa 7-Ketocholesterol-BSA using Mouse Anti-7-Ketocholesterol Monoclonal Antibody, Clone 7E1 (ASM10340). Lane 1: Molecular Weight Ladder (MW). Lane 2: BSA (0.5 μ g). Lane 3: BSA (2.0 μ g). Lane 4: 7-ketocholesterol-BSA (0.5 μ g). Lane 5: 7-ketocholesterol-BSA (2.0 μ g). Block: 5% Skim Milk in TBST. Primary Antibody: Mouse Anti-7-Ketocholesterol Monoclonal Antibody (ASM10340) at 1:1000 for 2 hours at RT. Secondary Antibody: Goat Anti-Mouse IgG: HRP at 1:2000 for 60 min at RT. Color Development: ECL solution for 5 min in RT. Predicted/Observed Size: 67 kDa.