

**SQSTM1 Antibody**  
**Catalog # ASM10518****Specification****SQSTM1 Antibody - Product Information**

|                   |                                |
|-------------------|--------------------------------|
| Application       | WB                             |
| Primary Accession | <a href="#">Q13501</a>         |
| Other Accession   | <a href="#">NP_001135770.1</a> |
| Host              | Rabbit                         |
| Reactivity        | Human                          |
| Clonality         | Polyclonal                     |

**Description**

Rabbit Anti-Human SQSTM1 Polyclonal

**Target/Specificity**

Detects ~60 kDa. Band at 40 kDa is isoform 2.

**Other Names**

EBI3-associated protein of 60 kDa Antibody, ORCA Antibody, Osi Antibody, SQSTM 1 Antibody, ZIP 3 Antibody, A170 Antibody, SQSTM1 Antibody, Ubiquitin binding protein p62 Antibody, Protein kinase C-zeta-interacting protein Antibody, FTDALS3 Antibody, Ubiquitin-binding protein p62 Antibody, PKC-zeta-interacting protein Antibody, EBIAP Antibody, Sqstm1 Antibody, Phosphotyrosine independent ligand for the Lck SH2 domain of 62 kDa Antibody, EBI3 associated protein p60 Antibody, EBI3 associated protein of 60 kDa Antibody, SQSTM\_HUMAN Antibody, OSF-6 Antibody, Paget disease of bone 3 Antibody, p62B Antibody, p60 Antibody, PDB 3 Antibody, EBI 3 associated protein p60 Antibody, MGC127197 Antibody, Phosphotyrosine-independent ligand for the Lck SH2 domain of 62 kDa Antibody, EBI 3 associated protein of 60 kDa Antibody, ZIP3 Antibody, STAP Antibody, p62 Antibody, Sequestosome 1 Antibody, OSIL Antibody, STONE14 Antibody, Phosphotyrosine independent ligand for the Lck SH2 domain p62 Antibody, PDB3 Antibody, Oxidative stress induced like Antibody, Sequestosome-1 Antibody, ZIP Antibody,

**Immunogen**

Synthetic peptide from the C-terminal of Human SQSTM1 (aa. 335-345)

**Purification**

Peptide Affinity Purified

**Storage**

-20°C

**Storage Buffer**

PBS, 50% glycerol, 0.09% sodium azide

**Shipping Temperature**

Blue Ice or 4°C

**Certificate of Analysis**

A 1:1000 dilution of SPC-637 was sufficient for detection of SQSTM1 in 15 µg of Human HeLa Cell Lysates by ECL immunoblot analysis using goat anti-rabbit IgG:HRP as the secondary antibody.

**Cellular Localization**


Cytoplasm | Late Endosome | Lysosome | Cytoplasmic Vesicle | Autophagosome | Nucleus | Endoplasmic Reticulum | Cytoplasm | P-Body

## SQSTM1 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


## SQSTM1 Antibody - Images



Western blot analysis of Human HeLa and 293Trap cell lysates showing detection of 47.7 kDa SQSTM1 protein using Rabbit Anti-SQSTM1 Polyclonal Antibody (ASM10518). Lane 1: Molecular Weight Ladder (MW). Lane 2: Human HeLa and 293Trap cell lysates. Load: 15 µg . Block: 5% Skim Milk in 1X TBST. Primary Antibody: Rabbit Anti-SQSTM1 Polyclonal Antibody (ASM10518) at 1:1000 for 1 hour at RT. Secondary Antibody: Goat Anti-Rabbit HRP at 1:2000 for 60 min at RT. Color Development: ECL solution for 6 min in RT. Predicted/Observed Size: 47.7 kDa. Other Band(s): ~60 in Jurkat.



Western blot analysis of Human Jurkat cell lysates showing detection of 47.7 kDa SQSTM1 protein using Rabbit Anti-SQSTM1 Polyclonal Antibody (ASM10518). Lane 1: Molecular Weight Ladder (MW). Lane 2: Human Jurkat cell lysates. Load: 15  $\mu$ g . Block: 5% Skim Milk in 1X TBST. Primary Antibody: Rabbit Anti-SQSTM1 Polyclonal Antibody (ASM10518) at 1:1000 for 1 hour at RT. Secondary Antibody: Goat Anti-Rabbit HRP at 1:2000 for 60 min at RT. Color Development: ECL solution for 6 min in RT. Predicted/Observed Size: 47.7 kDa. Other Band(s): ~60 .



Western blot analysis of Mouse Brain cell lysates showing detection of 47.7 kDa SQSTM1 protein using Rabbit Anti-SQSTM1 Polyclonal Antibody (ASM10518). Lane 1: Molecular Weight Ladder (MW). Lane 2: Mouse Brain cell lysates. Load: 15  $\mu$ g . Block: 5% Skim Milk in 1X TBST. Primary Antibody: Rabbit Anti-SQSTM1 Polyclonal Antibody (ASM10518) at 1:1000 for 1 hour at RT. Secondary Antibody: Goat Anti-Rabbit HRP at 1:2000 for 60 min at RT. Color Development: ECL solution for 6 min in RT. Predicted/Observed Size: 47.7 kDa.

#### SQSTM1 Antibody - Background

SQSTM1 or Sequestome 1 (p62) is a ubiquitin binding protein that interacts with ubiquitin, providing a scaffold for several signaling proteins and triggering degradation of proteins through the proteasome or lysosome. This gene regulates activation of the nuclear factor kappa-B (NF- $\kappa$ B) signaling pathway. The protein functions as a scaffolding/adaptor protein in concert with TNF receptor-associated factor 6 to mediate activation of NF- $\kappa$ B in response to upstream signals. Mutations in this gene result in sporadic and familial Paget disease of bone. SQSTM1 also binds

autophagosomal membrane protein LC3/ATG8.

#### **SQSTM1 Antibody - References**

1. Kirkin V., et al. (2009) Mol Cell. 34: 259-269.
2. Seibenhener M.L., et al. (2007) FEBS Lett. 581: 175-179.
3. Komatsu M., et al. (2010) Nat Cell Biol. 12: 213-223.
4. Bjørkøy G., et al. (2006) Autophagy. 2: 138-139.
5. Pankiv S., et al. (2007) J Biol Chem. 282: 24131-24145.